

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 296: 1-18

ISSN 1026-051X

July 2015

hppt/ urn:lsid:zoobank.org:pub: 892E2A5E-10C4-4BBD-A87B-C7F6CEC233BA

THE BEES OF THE GENUS *COLLETES* LATREILLE (HYMENOPTERA, COLLETIDAE) COLLECTED BY THE SOVIET-MONGOLIAN EXPEDITIONS 1967-1982

M. Yu. Proshchalykin¹⁾, M. Kuhlmann²⁾

1) Institute of Biology and Soil Science, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, 690022, Russia. E-mail: proshchalin@biosoil.ru

2) Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK. E-mail: m.kuhlmann@nhm.ac.uk

An annotated list of 26 species of *Colletes* collected in Mongolia by Soviet-Mongolian expeditions in 1967–1982 is given. Two species, *Colletes perforator* Smith, 1869 and *C. pauljohni* Kuhlmann, 2002 are newly recorded from Mongolia. An updated checklist of the 34 species of *Colletes* so far known from Mongolia is provided.

KEY WORDS. Bees, Apoidea, Apiformes, Palaearctic region, biodiversity, new records.

М. Ю. Прощалыкин¹⁾, М. Кульман²⁾. Пчелы рода *Colletes* Latreille (Hymenoptera, Halictidae), собранные в результате Советско-Монгольских экспедиций 1967-1982 гг. // Дальневосточный энтомолог. 2015. N 296. С. 1-18.

Приводится список 26 видов пчел из рода *Colletes*, собранных в Монголии в результате Советско-Монгольских экспедиций в 1967–1982 гг. и хранящихся в Зоологическом институте РАН, Санкт-Петербург. Из них два вида: *Colletes perforator* Smith, 1869 и *C. pauljohni* Kuhlmann, 2002 указываются впервые для

фауны Монголии. Дан уточненный список *Colletes* Монголии, насчитывающий 34 вида.

- 1) Биолого-почвенный институт ДВО РАН, Владивосток, 690022, Россия.
- 2) Музей естественной истории, Лондон, SW7 5BD, Великобритания.

INTRODUCTION

Mongolia is located in central Asia, bordered by Russia to the north and China in the south. It is the sixth largest country in Asia, covering 1,564,100 sq. km and is divided into 21 provinces called “aimags” (Fig. 1).



Fig. 1. Administrative map of Mongolia showing provinces (aimags) (from Kuhlmann & Proshchalykin, 2013a).

Soviet-Mongolian expeditions were conducted from 1967 to 1983 and collected extensive entomological material, which became the basis for the publication of numerous articles and books (including Insects of Mongolia in 11 volumes), devoted to the study of various insect families. However, taxonomic information about the Mongolian bee fauna is still fragmentary. Based on the material collected by these expeditions only four papers on Halictidae bees have been published (Pesenko, 1977, 1984a,b; Astafurova & Proshchalykin, 2015). In 2014 previously unstudied *Colletes* specimens collected by the Soviet-Mongolian expeditions were discovered in the Zoological Institute, Russian Academy of Sciences, St. Petersburg (ZISP) that complement previous studies (Kuhlmann & Dorn, 2002; Kuhlmann, 2009a; Kuhlmann & Proshchalykin, 2013a).

The results presented in this paper are based on 558 specimens collected by M.A. Kozlov (357 specimens), E.P. Narchuk (78), I.M. Kerzhner (63), E.S. Sugonyaev (44), V.F. Zaitsev (10), D. Myagmarsuren (3), E.L. Gur'eva (2), and G.S. Medvedev (1) in Mongolia during the eleven Soviet-Mongolian expeditions (1967–1971, 1973, 1975, 1976, 1978, 1980, 1982), that are currently housed in ZISP (Table 1).

Places and dates of collecting bees in Mongolia follows Emeljanov *et al.* (1968, 1973, 1977, 1978) and Kerzhner *et al.* (1982). Hard brackets are used when information is added to specimen label information (e.g. geographical coordinates) or published data (e.g. current name of an old place name; affiliation to a present day country).

In the present study we report additional records of 26 *Colletes* species with two species recorded from Mongolia for the first time. In total 34 *Colletes* species are now recorded from Mongolia (Table 2) with its centre of diversity in the South-West of the country (Fig. 2). *Colletes conradti* Noskiewicz, 1936 is removed from the list of Mongolian species (see Proshchalykin & Kuhlmann, 2015 for details).

We have used the following abbreviations for collectors: DM – D. Myagmarsuren, EG – E.L. Gur’eva, EN – E.P. Narchuk; ES – E.S. Sugonyaev; GM – G.S. Medvedev, IK – I.M. Kerzhner; MK – M.A. Kozlov. New distribution records are marked with an asterisk (*). The general distribution of *Colletes* follows Kuhlmann & Proshchalykin (2013a,b; 2014a,b), Proshchalykin & Kuhlmann (2015), and Niu *et al.* (2014). The definition of *Colletes* species-groups follows Noskiewicz (1936) and Kuhlmann *et al.* (2009).

Table 1. Details of the Soviet-Mongolian expeditions (1967–1982)

Years	Aimags	Collectors	Number of <i>Colletes</i> specimens/species
1967	Arkhangai, Bayan-Khongor, Uvurkhangai, Umnugovi, Ulaanbaatar, Tuv	I. Kerzhner, M. Kozlov, V. Zaitsev	15/11
1968	Usv, Khovd, Khuvsgul, Zavkhan	M. Kozlov	16/7
1969	Bayan-Khongor, Uvurkhangai, Umnugovi	I. Kerzhner, M. Kozlov, V. Zaitsev	43/7
1970	Usv, Khovd, Bayan-Khongor, Govi-Altai, Khuvsgul, Uvurkhangai, Selenge	I. Kerzhner, E. Narchuk, M. Kozlov, D. Myagmarsuren, V. Zaitsev	28/10
1971	Bayan-Khongor, Khentii, Dornogovi, Dornod, Sukhbaatar	I. Kerzhner, M. Kozlov	35/5
1973	Umnugovi	G. Medvedev	1/1
1975	Khuvsgul, Arkhangai, Selenge, Khentii, Dornogovi, Dornod, Sukhbaatar	E. Narchuk, E. Sugonyaev, M. Kozlov	177/18
1976	Tuv, Khentii, Dornod, Sukhbaatar	I. Kerzhner, M. Kozlov	165/9
1978	Bayan-Ulgii, Uvs, Bayan-Khongor, Govi-Altai	M. Kozlov, E. Gur’eva	28/8
1980	Khovd, Govi-Altai, Arkhangai	I. Kerzhner, M. Kozlov	39/8
1982	Bayan-Khongor	I. Kerzhner	1/1
		Total:	558/26

Table 2. The list of *Colletes* bees of Mongolia

№	Species	Reference date			
		I	II	III	IV
1	<i>C. alini</i> Kuhlmann, 2000	+	+	+	+
2	<i>C. arsenjevi</i> Kuhlmann, 2006		+	+	+
3	<i>C. chengtehensis</i> Yasumatsu, 1935	+	+	+	+
4	<i>C. cinerascens</i> Morawitz, 1893	+	+	+	+
5	<i>C. collaris</i> Dours, 1872	+	+	+	+
6	<i>C. cunicularius</i> (Linnaeus, 1761)	+		+	
7	<i>C. daviesanus</i> Smith, 1846	+	+	+	+
8	<i>C. dorni</i> Kuhlmann, 2002	+		+	
9	<i>C. dubitatus</i> Noskiewicz, 1936			+	+
10	<i>C. ebmeri</i> Kuhlmann, 2002	+	+	+	+
11	<i>C. edentuloides</i> Kuhlmann, 2011			+	
12	<i>C. edentulus</i> Noskiewicz, 1936	+	+	+	
13	<i>C. emaceatus</i> Noskiewicz, 1936		+	+	+
14	<i>C. floralis</i> Eversmann, 1852	+	+	+	+
15	<i>C. fodiens</i> (Fourcroy, 1785)	+		+	+
16	<i>C. fulvicornis</i> Noskiewicz, 1936	+	+	+	+
17	<i>C. gusi</i> Kuhlmann, 2009		+	+	+
18	<i>C. impunctatus</i> Nylander, 1852	+	+	+	+
19	<i>C. inexpectatus</i> Noskiewicz, 1936	+		+	
20	<i>C. jankowskyi</i> Radoszkowski, 1891	+	+	+	+
21	<i>C. kaszabi</i> Kuhlmann, 2002	+	+	+	+
22	<i>C. kozlovi</i> Friese, 1913	+	+	+	+
23	<i>C. mixtus</i> Radoszkowski, 1891	+		+	+
24	<i>C. pauljohni</i> Kuhlmann, 2002				+
25	<i>C. perforator</i> Smith, 1869				+
26	<i>C. plumuloides</i> Kuhlmann et Proshchalykin, 2013			+	
27	<i>C. popovi</i> Noskiewicz, 1936			+	+
28	<i>C. pseudocinerascens</i> Noskiewicz, 1936	+	+	+	+
29	<i>C. ravulus</i> Noskiewicz, 1936	+		+	+
30	<i>C. reinigi</i> Noskiewicz, 1936		+	+	
31	<i>C. roborovskyi</i> Friese, 1913	+		+	
32	<i>C. sidemii</i> Radoszkowski, 1891	+	+	+	+
33	<i>C. wacki</i> Kuhlmann, 2002	+	+	+	+
34	<i>C. wahrmani</i> Noskiewicz, 1959		+	+	+
<i>Total:</i>		23	21	32	26

References data source: I – Kuhlmann & Dorn, 2002; II – Kuhlmann, 2009; III – Kuhlmann & Proshchalykin, 2013; IV – current data.




Fig. 2. Number of Mongolian *Colletes* species by aimags (the names of the aimags in Fig. 1). Number of *Colletes* species recorded from Mongolian aimags/number of *Colletes* species collected by Soviet-Mongolian expeditions (1967–1982).

LIST OF SPECIES

Colletes caspicus species-group

Colletes alini Kuhlmann, 2000

Colletes alini Kuhlmann, 2000: 161–162, ♀, ♂ (type locality: “Mandschurei, Charbin”, Heilongjiang, China); Kuhlmann & Dorn, 2002: 90; Kuhlmann, 2009a: 21; Kuhlmann & Proshchalykin, 2011: 7; 2013a: 256; Proshchalykin, 2012: 449; Niu *et al.*, 2014: 455; Proshchalykin & Kuhlmann, 2015: 336.

Colletes seitzi: Osytshnuk & Romankova, 1995: 483; Proshchalykin, 2007: 879.

SPECIMENS EXAMINED. Mongolia. *Uvs Aimag*: 50 km E Ulangom [92°70'E 49°59'N], 10-11.VII 1968, 3 ♂ (MK); Togtokhyn-Shil Ridge, 50 km ESE Ulangom [92°04'E 49°59'N], 7.VIII 1970, 2 ♂ (IK); *Bayan-Khongor Aimag*: Orog-Noor Lake [100°42'E 45°03'N], 15-16.VIII 1967, ♂ (VZ); Shine-Dzhinst, Uluziin-Gol River [99°05'E 44°25'N], 15.VII 1982, 1 ♂ (IK); *Selenge Aimag*: 25 km E Darkhan, Sharyn-Gol Mt. [106°15'E 49°10'N], 30.VII 1975, ♂ (MK); *Tuv Aimag*: Kerulen, Kerulen River [108°30'E 47°50'N], 15.VIII 1976, ♀ (MK); *Khentii Aimag*: 25 km NE Bayan-Adraga [111°50'E 48°30'N], 6-7.VII 1976, 7 ♂ (IK); *Dornogovi Aimag*: 60 km NW Choir, Sumber [108°45'E 46°35'N], 12.VIII 1975, ♂ (MK); 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, ♂ (EN); 70 km ENE Sain-Shand, Dolotyn-Khuduk [111°10'E 44°45'N], 17.VIII 1975, ♂ (EN); *Dornod Aimag*: 7 km S Erentsav [114°80'E 49°61'N], 22.VIII 1975, ♂ (EN); 60 km ENE Bayan-Burd, Derkhin-Tsagan-Obo Mt. [119°00'E 46°80'N], 3-4.VII 1976, 2 ♂ (MK); 33 km SE Khalkh-Gol, Khalkin-Gol River [118°25'E 47°90'N], 31.VII 1976, 8 ♂ (MK); *Sukhbaatar Aimag*: 80 km NNE Barun-Urt [113°70'E 47°10'N], 19.VIII 1975, ♂ (EN); Dzotol-Khan-Ula Mt. [114°00'E 45°33'N], 20.VII 1976, ♀ (MK).

DISTRIBUTION. Mongolia (Uvs Aimag, Khovd Aimag, Bayan-Khongor Aimag, Umnugovi Aimag, *Selenge Aimag, Tuv Aimag, *Khentii Aimag, *Dornogovi Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (Eastern Siberia, Far East), China (Beijing, Tianjin, Hebei, Inner Mongolia, Heilongjiang, Liaoning, Shandong, Shanxi, Jiangsu, Ningxia, Xinjiang).

Colletes squamosus species-group

Colletes wahrmani Noskiewicz, 1959

Colletes wahrmani Noskiewicz, 1959: 43–45, ♂ (type locality: Bevsekhir, Turkey); Kuhlmann, 2009a: 21; 2009b: 16; Kuhlmann & Proshchalykin, 2013a: 257; 2013b: 416; Niu *et al.*, 2014: 472.

SPECIMENS EXAMINED. **Mongolia.** *Umnugovi Aimag:* 60 km E Talyn-Bilgakh-Bulak River [104°47'E 44°12'N], 17-19.VIII 1969, ♂ (IK); *Dornogovi Aimag:* 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, ♀ (EN).

DISTRIBUTION. Mongolia (Govi-Alтай Aimag, Umnugovi Aimag, *Dornogovi Aimag), China (Xinjiang), Kazakhstan, Uzbekistan, Turkmenistan, Pakistan, Turkey, Armenia.

Colletes mixtus species-group

Colletes dubitatus Noskiewicz, 1936

Colletes dubitatus Noskiewicz, 1936: 241–242, ♂ (type locality: Tartugai, Kazakhstan); Kuhlmann & Proshchalykin, 2013a: 257; 2013b: 417.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag:* 12 km S Altai [92°22'E 45°83'N], 22.VII 1970, ♀ (IK); *Bayan-Khongor Aimag:* Ekhin-Gool Oasis [99°81'E 42°31'N], 11-14.VIII 1969, ♀, 3 ♂ (MK).

DISTRIBUTION. Mongolia (Khovd Aimag, Bayan-Khongor Aimag), Kazakhstan, Uzbekistan, Turkmenistan, Tajikistan.

Colletes kozlovi Friese, 1913

Colletes kozlovi Friese, 1913: 59, ♀, ♂ (type locality: “Mongolia” [China: Xinjiang]); Noskiewicz, 1936: 240; Kuhlmann & Dorn, 2002: 91; Kuhlmann, 2005: 1369; 2009a: 21; 2009b: 16; Kuhlmann & Proshchalykin, 2011: 7; 2013a: 257; 2013b: 418; Niu *et al.*, 2014: 470; Proshchalykin & Kuhlmann, 2015: 336.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag:* 50 km SSW Uench, Utyn-Mod [91°45'E 45°55'N], 27.VI 1980, 2 ♀, 12 ♂ (MK); *Govi-Alтай Aimag:* 20 km ENE Khatan-Khairkhan-Ula Mt. [96°30'E 45°00'N], 4-5.VIII 1978, 5 ♂ (MK); 5.VIII 1978, ♂ (EG); *Arkhangai Aimag:* Terkhin-Tsaagan-Nuur Lake, 15 km W Taryat [99°43'E 48°09'N], 19.VII 1980, ♂ (MK); *Dornogovi Aimag:* 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, 2 ♀ (MK); 14.VIII 1975, ♂ (EN).

DISTRIBUTION. Mongolia (Khovd Aimag, Zavkhan Aimag, *Govi-Altai Aimag, *Arkhangai Aimag, Bayan-Khongor Aimag, Umnugovi Aimag, Dornogovi Aimag, Dornod Aimag), Russia (South-East of European part, Eastern Siberia), China (Inner Mongolia, Qinghai, Gansu, Ningxia, Xinjiang), Kazakhstan, Uzbekistan, Kyrgyzstan, Turkmenistan, Tajikistan, Azerbaijan.

***Colletes mixtus* Radoszkowski, 1891**

Colletes mixtus Radoszkowski, 1891: 257–258, ♀, ♂ (type locality: no locality); Kuhlmann & Dorn, 2002: 91; Kuhlmann, 2005: 1370; Kuhlmann & Proshchalykin, 2013a: 258; Niu et al., 2014: 471.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag:* 25 km S Altai, Elkhon [92°20'E 45°36'N], 22-23.VI 1980, 1♀, 4♂ (MK); *Bayan-Khongor Aimag:* Ekhin-Gool Oasis [99°81'E 42°31'N], 11-14.VIII 1969, 3♀, 19♂ (MK); 31.VIII 1970, ♂ (MK); Talyn-Bilgekh-Bulak, 13 km E Tsagan-Bogdo-Ula [99°23'E 42°85'N], 16.VIII 1969, 4♀, 6♂ (MK); 100 km SE Bayan-Obo [101°12'E 45°10'N], 21.VI 1971, ♂ (MK).

DISTRIBUTION. Mongolia (Khovd Aimag, Govi-Altai Aimag, Bayan-Khongor Aimag), China (Gansu, Xinjiang), Kazakhstan, Uzbekistan, Turkmenistan, Tajikistan, Turkey.

***Colletes roborovskyi* species-group**

***Colletes roborovskyi* Friese, 1913**

Colletes roborovskyi Friese, 1913: 59, ♀, ♂ (type locality: "Mongolia" [China: Xinjiang]); Kuhlmann & Proshchalykin, 2013a: 258; 2013b: 419.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag:* 25 km N Bulgan, Ulyastain-Gol River [91°85'E 46°32'N], 31.VII 1970, 4♂ (MK).

DISTRIBUTION. Mongolia (Khovd Aimag, Umnugovi Aimag), Russia (South-East of European part, Eastern Siberia), China (Inner Mongolia, Xinjiang), Kazakhstan, Kyrgyzstan, Turkmenistan, Tajikistan, Iran, Pakistan, Turkey, Georgia, Azerbaijan.

***Colletes uralensis* species-group**

***Colletes kaszabi* Kuhlmann, 2002**

Colletes kaszabi Kuhlmann in Kuhlmann & Dorn, 2002: 92, ♀, ♂ (holotype: ♀, "Menengijntal 80 km WSW vom SW Ecke des Sees Bujr nur", Dornod Aimag, Mongolia); Kuhlmann, 2009a: 21; Kuhlmann & Proshchalykin, 2013a: 258; Niu et al., 2014: 473; Proshchalykin & Kuhlmann, 2015: 338.

SPECIMENS EXAMINED. **Mongolia.** *Govi-Altai Aimag:* 20 km ENE Khatan-Khairkhan-Ula Mt. [96°30'E 45°00'N], 4-5.VIII 1978, 5♂ (MK); 5.VIII 1978, ♂

(EG); 25 km NE Bayan [96°00'E 46°30'N], 16.VI 1980, 2 ♂ (MK); *Khuvsgul Aimag*: 20 km SE Toson-Tsengel, Selenga River [100°50'E 49°25'N], 25.VII 1975, ♂ (ES); *Tuv Aimag*: Kerulen, Kerulen River [108°30'E 47°50'N], 15.VIII 1976, ♀ (MK); *Dornogovi Aimag*: 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, 2 ♀, 16 ♂ (EN); 10 km E Dzun-Bayan [110°03'E 44°18'N], 15.VIII 1975, 5 ♂ (ES); 70 km ENE Sain-Shand, Dolotyn-Khuduk [111°10'E 44°45'N] 17.VIII 1975, ♂ (EN); *Sukhbaatar Aimag*: 15 km SE Khongor, Ongon-Els [113°09'E 45°72'N], 5-6.VII 1971, ♂ (MK); Dzotol-Khan-Ula Mt. [114°00'E 45°33'N], 20.VII 1976, ♂ (MK).

DISTRIBUTION. Mongolia (Uvs Aimag, Zavkhan Aimag, Govi-Altai Aimag, *Khuvsgul Aimag, Bayan-Khongor Aimag, Bulgan Aimag, Ulaanbaatar, Tuv Aimag, Dornogovi Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (Eastern Siberia) China (Inner Mongolia).

Colletes flavicornis species-group

Colletes emaceatus Noskiewicz, 1936

Colletes emaceatus Noskiewicz, 1936: 264–267, ♂ (type locality: Muin-Kum desert, Kazakhstan); Kuhlmann, 2009a: 22; Kuhlmann & Proshchalykin, 2013a: 258; 2013b: 420.

SPECIMENS EXAMINED. **Mongolia.** *Bayan-Khongor Aimag*: Tuin-Gool River, S Bogdo-Ula [101°04'E 45°09'N], 5.VIII 1969, ♂ (IK); *Umnugovi Aimag*: Bordzon-Goby, 80 km SSE Nomgon [105°77'E 42°35'N], 5-8.VIII 1967, ♀, ♂ (IK); *Dornogovi Aimag*: 70 km ENE Sain-Shand, Dolotyn-Khuduk [111°10'E 44°45'N], 17.VIII 1975, 4 ♀ (EN); 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, 2 ♀ (MK); 15 km E Dzun-Bayan [110°12'E 44°26'N], 15.VIII 1975, ♂ (ES).

DISTRIBUTION. Mongolia (Bayan-Khongor Aimag, Uvurkhangai Aimag, Umnugovi Aimag, *Dornogovi Aimag), Kazakhstan, Turkmenistan.

Colletes gusi Kuhlmann, 2009

Colletes gusi Kuhlmann, 2009a: 22, ♀, ♂ (holotype: ♂, 28 km SE Chatun-Bulag, Dornogovi Aimag, Mongolia); Kuhlmann & Proshchalykin, 2013a: 259.

SPECIMENS EXAMINED. **Mongolia.** *Usv Aimag*: 50 km E Ulangoma, Uvs-Noor Lake [92°15'E 50°22'N], 6.VIII 1970, ♂ (MK); *Umnugovi Aimag*: 60 km E Talyn-Bilgakh-Bulak River [104°45'E 44°06'N], 14-19.VIII 1969, ♀ (MK); *Dornogovi Aimag*: 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, ♀ (EN); 14.VIII 1975, ♂ (MK).

DISTRIBUTION. Mongolia (Usv Aimag, Umnugovi Aimag, Dornogovi Aimag, Dornod Aimag).

Colletes popovi Noskiewicz, 1936

Colletes popovi Noskiewicz, 1936: 271–274, ♀ (type locality: “Schibendu Schigusa s. Satschjou, Gasch. Gob.” [China: Xinjiang]); Kuhlmann, 2005: 1372; Kuhlmann & Proshchalykin, 2013a: 259; 2013b: 423.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag*: 12 km SW Altaj-som, Bodonchin-Gol River [92°25'E 45°81'N], 22.VII 1970, ♂ (MK); *Govi-Altai Aimag*: 60 km SW Tsel [95°08'E 45°42'N], 19.VII 1970, ♂ (VZ); *Bayan-Khongor Aimag*: Talyn-Bilgekh-Bulak, 13 km E Tsagan-Bogdo-Ula [99°23'E 42°85'N], 16.VIII 1969, ♂ (MK); *Uvurkhangai Aimag*: Tatsyn-Tsagan-Noor Lake [101°61'E 44°85'N], 2-4.VIII 1969, ♀ (VZ).

DISTRIBUTION. Mongolia (Khovd Aimag, Govi-Altai Aimag, Bayan-Khongor Aimag, Uvurkhangai Aimag), China (Qinghai), Kazakhstan.

Colletes fodiens species-group

Colletes daviesanus Smith, 1846

Colletes daviesanus Smith, 1846: 1278, ♀, ♂ (type locality: England); Kuhlmann & Dorn, 2002: 93; Kuhlmann, 2005: 1373; 2009a: 24; Proshchalykin & Kuhlmann, 2012: 15; 2015: 330; Kuhlmann & Proshchalykin, 2013a: 260; 2013b: 424; Proshchalykin, 2012: 449; Niu *et al.*, 2014: 457.

SPECIMENS EXAMINED. **Mongolia.** *Umnugovi Aimag*: 65 km SE Nomgon [105°91'E 42°35'N], 9.VIII 1967, ♂ (IK).

DISTRIBUTION. Mongolia (Uvs Aimag, Zavkhan Aimag, Umnugovi Aimag, Tuv Aimag), Russia (European part, Siberia, Far East), China (Qinghai, Heilonjiang), Kazakhstan, Kyrgyzstan, Uzbekistan, Europe, Turkey, Armenia, Georgia.

Colletes pauljohni Kuhlmann, 2002

Colletes pauljohni Kuhlmann, 2002: 750-751, ♀, ♂ (type locality: south shore of Issyk-Kul Lake, Teploklyuchenka [Ak-su], Kyrgyzstan).

SPECIMENS EXAMINED. **Mongolia.** *Bayan-Khongor Aimag*: 10 km ENE Samon-Khairkhan-Ula Mt. [97°60'E 44°35'N], 9-10.VIII 1978, ♀, ♂ (MK).

DISTRIBUTION. *Mongolia (*Bayan-Khongor Aimag), China (Xinjiang), Kazakhstan, Kyrgyzstan, Tajikistan.

Colletes perforator Smith, 1869

Colletes perforator Smith, 1869: 205, ♀ (type locality: Japan).

SPECIMENS EXAMINED. **Mongolia.** *Selenge Aimag*: Dulan-Khan, Ero-Gol River [106°E 30' 49°56'N], 4.VIII 1975, 2 ♀ (EN); 4.VIII 1975, ♂ (MK).

DISTRIBUTION. *Mongolia (*Selenge Aimag), Russia (Eastern Siberia, Far East), China (Beijing, Shanxi, Inner Mongolia, Heilongjiang, Jilin, Jiangxi), Japan (Hokkaido, Honshu, Shikoku, Kyushu, Tsushima).

Colletes ravulus Noskiewicz, 1936

Colletes ravulus Noskiewicz, 1936: 335-337, ♂ (type locality: "Nordmongolei", Mongolia); Kuhlmann & Dorn, 2002: 94; Kuhlmann & Proshchalykin, 2011: 10; 2013a: 260; Niu *et al.*, 2014: 461; Proshchalykin & Kuhlmann, 2015: 336.

SPECIMENS EXAMINED. **Mongolia.** *Usv Aimag*: channel between Airag and Khirgis-Noor [93°20'E 49°10'N], 28-29.VIII 1968, 2 ♀ (MK); *Dornod Aimag*: 13 km W Dash-Balbar, Uldza River [114°00'E 49°20'N], 24.VIII 1975, 2 ♀ (EN).

DISTRIBUTION. Mongolia (Usv Aimag, Dornod Aimag), Russia (Eastern Siberia), China (Xinjiang, Shanxi, Inner Mongolia).

Colletes clypearis species-group

Colletes cinerascens Morawitz, 1893

Colletes cinerascens Morawitz, 1893: 80, ♂ (type locality: “Jagnob, Kol”, Tajikistan); Kuhlmann & Dorn, 2002: 95; Kuhlmann, 2005: 1377; 2009a: 25; 2009b: 18; Kuhlmann & Proshchalykin, 2013a: 261; 2013b: 429; Niu *et al.*, 2014: 474; Proshchalykin & Kuhlmann, 2015: 332.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag*: 20 km N Bulgan, Ulyastain-Gol River [92°10'E 46°05'N], 30.VI-1.VII 1980, ♂ (MK); *Bayan-Ulgii Aimag*: 20 km NW Bulgan, Ikh-Dzhargalantyn-Gol River [91°00'E 47°00'N], 22-23.VII 1978, ♂ (MK).

DISTRIBUTION. Mongolia (*Bayan-Ulgii Aimag, Uvs Aimag, Khovd Aimag, Govi-Altai Aimag, Bayan-Khongor Aimag), Russia (Eastern Siberia), China (Gansu, Qinghai, Xinjiang), Kazakhstan, Kyrgyzstan, Tajikistan.

Colletes ebmeri Kuhlmann, 2002

Colletes ebmeri Kuhlmann in Kuhlmann & Dorn, 2002: 95, ♀, ♂ (holotype: ♂, “18 km S Pass Egijn davaa”, Bayan-Khongor Aimag, Mongolia); Kuhlmann, 2009a: 25; Kuhlmann & Proshchalykin, 2013a: 261; 2013b: 430; Proshchalykin & Kuhlmann, 2015: 332.

SPECIMENS EXAMINED. **Mongolia.** *Khovd Aimag*: 40 km N Uench [92°06'E 46°38'N], 31.VII-1.VIII 1968, ♀ (MK); 15 km WNW Dut [91°30'E 47°37'N], 8.VII 1980, ♀, 2 ♂ (MK); *Govi-Altai Aimag*: 37 km NE Tsogt, Dutiin-Daba Ridge [97°13'E 45°67'N], 14.VII 1970, ♂ (MK); *Khuvsgul Aimag*: 20 km SW Buren-Khan [99°85'E 49°51'N], 30.VI 1968, ♂ (MK); 25 km km NNE Ulan-Ula [99°13'E 50°40'N], 15.VII 1975, 2 ♀, ♂ (EN); 17 km N Shine-Ider [99°50'E 49°10'N], 21.VII 1975, ♀ (ES); *Ulaanbaatar* [106°59'E 47°45'N], 11.VI 1967, ♂ (IK); *Dornogovi Aimag*: 60 km NW Choir, Sumber [108°45'E 46°35'N], 12.VIII 1975, 3 ♂ (MK).

DISTRIBUTION. Mongolia (Khovd Aimag, Govi-Altai Aimag, Khuvsgul Aimag, Arkhangai Aimag, Bayan-Khongor Aimag, Ulaanbaatar, *Dornogovi Aimag), Russia (Eastern Siberia), Kazakhstan.

Colletes floralis Eversmann, 1852

Colletes floralis Eversmann, 1852: 46, ♀, ♂ (type locality: “in prov. Orenburg., Saratov. et Astrachanensi”, Russia); Kuhlmann & Dorn, 2002: 96; Kuhlmann, 2005: 1377; 2009a: 28; 2009b: 18; Proshchalykin, 2007: 878; 2012: 449; Kuhlmann & Proshchalykin, 2011: 15; 2013a: 261; 2013b: 430; Niu *et al.*, 2014: 475; Proshchalykin & Kuhlmann, 2015: 332.

SPECIMENS EXAMINED. **Mongolia.** *Bayan-Ulgii Aimag*: 20 km NW Bulgan, Ikh-Dzhargalantyn-Gol River [91°00'E 47°00'N], 22-23.VII 1978, ♂ (MK); *Selenge Aimag*: 25 km E Darkhan, Sharyn-Gol Mt. [106°15'E 49°10'N], 30.VII 1975, ♀ (MK); *Ulaanbaatar* [106°59'E 47°45'N], 14.VII 1967, ♂ (VZ); *Tuv Aimag*: Kerulen [108°30'E 47°50'N], 15.VIII 1976, 2 ♂ (MK); *Dornod Aimag*: 13 km W Dash-Balbar, Uldza River [114°00'E 49°20'N], 24.VIII 1975, ♀ (EN); 32 km SE Salkhit Mt., Numregin-Gol River [119°45'E 47°00'N], 9.VIII 1976, ♂ (MK); *Sukhbaatar Aimag*: 65 km NNW Dariganga [113°10'E 45°55'N], 14.VII 1976, 2 ♂ (MK).

DISTRIBUTION. Mongolia (**Bayan-Ulgii Aimag*, *Khuvsgul Aimag*, *Arkhangai Aimag*, *Bayan-Khongor Aimag*, *Bulgan Aimag*, *Uvurkhangai Aimag*, *Selenge Aimag*, *Ulaanbaatar*, *Tuv Aimag*, **Dornod Aimag*, **Sukhbaatar Aimag*), Russia (European part, Siberia, Far East), China (Hebei, Qinghai, Xinjiang, Xizang, Guangdong, Yunnan), Japan (Hokkaido, Honshu), Kazakhstan, Uzbekistan, Kyrgyzstan, Tajikistan, Iran, Europe, Georgia, India.

Colletes fulvicornis Noskiewicz, 1936

Colletes fulvicornis Noskiewicz, 1936: 416, ♂ (lectotype: ♂, designated by Kuhlmann, 2000: 180, Cholt, Gobi, 16.VI 1926, Kozlov leg., Mongolia); Osytshnjuk & Romankova, 1995: 483; Kuhlmann & Dorn, 2002: 96; Kuhlmann, 2009a: 28; Kuhlmann & Proshchalykin, 2011: 16; 2013a: 262; Proshchalykin & Kuhlmann, 2015: 333.

SPECIMENS EXAMINED. **Mongolia.** *Uvs Aimag*: 20 km S Under-Khangai, Chigdzhii-Gol River [95°00'E 48°30'N], 2.VII 1978, 2 ♀, 3 ♂ (MK); *Khuvsgul Aimag*: 25 km NNE Ulan-Ula, Shikshid-Gol River [99°13'E 50°40'N], 15.VII 1975, ♂ (EN); *Umnugovi Aimag*: 40 km S Bulgan, Gurvan-Saikhan Ridge [103°87'E 43°78'N], 28-29.VII 1967, ♀ (IK); *Khentii Aimag*: 12 km N Gal-Shara [111°08'E 46°84'N], 30.VII 1971, ♂ (MK); *Dornod Aimag*: Khukh-Nur Lake [115°40'E 49°35'N], 25.VI 1976, ♂ (MK); *Sukhbaatar Aimag*: Dzotol-Khan-Ula Mt. [114°00'E 45°33'N], 20.VII 1976, 8 ♀, 26 ♂ (MK).

DISTRIBUTION. Mongolia (*Uvs Aimag*, *Khovd Aimag*, *Govi-Altai Aimag*, **Khuvsgul Aimag*, *Arkhangai Aimag*, *Bayan-Khongor Aimag*, *Uvurkhangai Aimag*, *Umnugovi Aimag*, *Ulaanbaatar*, *Tuv Aimag*, *Dundgovi Aimag*, *Khentii Aimag*, *Dornod Aimag*, **Sukhbaatar Aimag*), Russia (Siberia), China (Xinjiang, Xizang).

Colletes impunctatus Nylander, 1852

Colletes impunctatus Nylander, 1852: 249, ♀, ♂ (type locality: North Sweden); Kuhlmann & Dorn, 2002: 97; Proshchalykin, 2007: 879; 2012: 450; Kuhlmann, 2009a: 28; Kuhlmann & Proshchalykin, 2011: 17; 2013a: 263; 2013b: 431; Proshchalykin & Kuhlmann, 2015: 333.

SPECIMENS EXAMINED. **Mongolia.** *Bayan-Ulgii Aimag*: 20 km NW Bulgan, Ikh-Dzhargalantyn-Gol River [91°00'E 47°00'N], 22-23.VII 1978, ♂ (MK); *Uvs Aimag*: 20 km S Under-Khangai, Chigdzhii-Gol River [95°00'E 48°30'N], 2.VII 1978, ♀ (MK); Ureg-Nur Lake, mouth of Karga River [91°00'E 50°20'N],

6.VII 1978, ♀ (MK); *Khovd Aimag*: 40 km N Uench-Somon [92°06'E 46°38'N], 31.VII-1.VIII 1968, ♂ (MK); *Zavkhan Aimag*: 30 km WNW Tes-Somon [96°13'E 48°25'N], 3-4.VII 1968, ♂ (MK); *Govi-Altai Aimag*: 8 km NE Argalant-Ula Mt. [93°05'E 45°50'N], 20.VI 1980, ♂ (MK); 25 km NE Bayan [96°00'E 46°30'N], 16.VI 1980, 2 ♂ (MK); *Khuvsugul Aimag*: Delger-Muren River, Buren-Khan [99°02'E 50°21'N], 28-29.VI 1968, 2 ♂ (MK); 25 km NNE Ulan-Ula, Shikshid-Gol River [99°13'E 50°40'N], 15.VII 1975, 4 ♀, 6 ♂ (EN); 15.VII 1975, 2 f, ♂ (ES); 17 km N Shine-Ider [99°50'E 49°10'N], 22.VII 1975, 2 ♀ (EN); 22.VII 1975, 6 ♀, 6 ♂ (ES); 20 km SE Toson-Tsengel, Selenga River [100°50'E 49°25'N], 25.VII 1975, ♂ (EN); *Arkhangai Aimag*: Terkhin-Tsaagan-Nuur Lake, 15 km W Taryat [99°43'E 48°09'N], 19.VII 1980, ♀, ♂ (MK); Chulutyn-Gol River, Chulut [100°18'E 47°65'N], 29.VI 1975, 7 ♀, 4 ♂ (MK); *Tuv Aimag*: Mungen-Mor't [108°42'E 48°17'N], 6-7.VIII 1967, 2 ♂ (MK); *Ulaanbaatar*, Tola River [106°55'E 47°55'N], 19.VI 1967, ♀ (IK); *Dornod Aimag*: Khukh-Nur Lake [115°40'E 49°35'N] 25.VI 1976, 2 ♂ (MK).

DISTRIBUTION. Mongolia (Bayan-Ulgii Aimag, Uvs Aimag, Khovd Aimag, Zavkhan Aimag, Govi-Altai Aimag, Khuvsugul Aimag, Arkhangai Aimag, Bayan-Khongor Aimag, Uvurkhangai Aimag, Umnugovi Aimag, Ulaanbaatar, Tuv Aimag, Dornod Aimag), Russia (European part, Siberia, Far East), China (Inner Mongolia), Kazakhstan, Europe.

Colletes jankowskyi Radoszkowski, 1891

Colletes jankowskyi Radoszkowski, 1891: 253, ♀ (type locality: “sur les bords de la riviere Sidémi (non loin de Vladivostok)”, Primorskiy Terr., Russia); Osytshnjuk & Romankova, 1995: 484; Kuhlmann & Dorn, 2002: 98; Kuhlmann, 2009a: 29; 2009b: 18; Kuhlmann & Proshchalykin, 2011: 17; 2013a: 263; 2013b: 431; Proshchalykin, 2012: 450; Proshchalykin & Kuhlmann, 2015: 334.

SPECIMENS EXAMINED. **Mongolia**. *Uvurkhangai Aimag*: Arts-Bogdo Ridge, 20 km S Khovd [102°15'E 44°43'N], 12-13.VIII 1967, ♀ (VZ); *Khentii Aimag*: 8 km S Norovlin [111°59'E 48°40'N], 25.VIII 1975, ♂ (MK); *Dornod Aimag*: 13 km W Dash-Balbar, Uldza River [114°00'E 49°20'N], 24.VIII 1975, 2 ♀ (EN).

DISTRIBUTION. Mongolia (Uvs Aimag, Arkhangai Aimag, Bayan-Khongor Aimag, Uvurkhangai Aimag, Ulaanbaatar, Tuv Aimag, *Khentii Aimag), Russia (Siberia, Far East), China (Beijing, Hebei, Shandong, Shanxi, Gansu, Heilongjiang, Jilin, Inner Mongolia, Yunnan), Tajikistan.

Colletes pseudocinerascens Noskiewicz, 1936

Colletes pseudocinerascens Noskiewicz, 1936: 424, ♀, ♂ (lectotype: ♂, designated by Kuhlmann, 2000: 181, Jakutsk, Russia); Kuhlmann & Dorn, 2002: 98; Kuhlmann, 2009a: 29; Kuhlmann & Proshchalykin, 2011: 18; 2013a: 263; Proshchalykin & Kuhlmann, 2015: 334.

SPECIMENS EXAMINED. **Mongolia**. *Usv Aimag*: Togtokhyn-Shil Ridge, 50 km ESE Ulangom [92°04'E 49°59'N], 7.VIII 1970, 3 ♀ (MK); *Zavkhan Aimag*:

30 km WNW Tes-Somon [96°13'E 48°25'N], 3-4.VII 1968, ♂ (MK); *Khuvsgul Aimag*: 25 km NNE Ulan-Ula [99°13'E 50°40'N], 15-16.VII 1975, 2 ♀, ♂ (MK).

DISTRIBUTION. Mongolia (Usv Aimag, Zavkhan Aimag, Khuvsgul Aimag), Russia (Siberia).

***Colletes sidemii* Radoszkowski, 1891**

Colletes sidemii Radoszkowski, 1891: 252 (type locality: “sur les bords de la riviere Sidémi (non loin de Vladivostok)”, Primorskiy Terr., Russia); Osytshnjuk & Romankova, 1995: 484; Kuhlmann & Dorn, 2002: 98; Proshchalykin, 2007: 879; 2012: 450; Kuhlmann, 2005: 1378; 2009a: 29; Kuhlmann & Proshchalykin, 2011: 19; 2013a: 264; 2013b: 431; Niu *et al.*, 2014: 477; Proshchalykin & Kuhlmann, 2015: 335.

SPECIMENS EXAMINED. **Mongolia.** *Usv Aimag*: 5 km S Khyargas-Noor [93°20'E 49°10'N], 10.VIII 1970, ♂ (VZ); 20 km S Under-Khangai, Chigdziin-Gol River [95°00'E 48°30'N], 2.VII 1978, ♀, 3 ♂ (MK); *Khuvsgul Aimag*: Delger-Muren River, Buren-Khan [99°02'E 50°21'N], 28-29.VI 1968, ♂ (MK); 17 km N Shine-Ider [99°50'E 49°10'N], 22.VII 1975, ♀, ♂ (EN); 20 km SE Toson-Tsengel, Selenga River [100°50'E 49°25'N], 25.VII 1975, ♂ (ES); Ider River, Dzhargalant [100°83'E 48°79'N], 19-20.VII 1975, 2 ♂ (MK); *Arkhangai Aimag*: 30 km E Tsetserleg [102°00'E 47°20'N], 1.VII 1975, ♂ (MK); 2.VII 1975, ♂ (ES); *Uvurkhangai Aimag*: Arts Bogdo Ridge [102°08'E 44°65'N], 11.VII 1970, 3 ♂ (DM); *Selenge Aimag*: 25 km E Darkhan, Sharyn-Gol Mt. [106°15'E 49°10'N], 30.VII 1975, 4 ♀, 4 ♂ (MK); 30.VII 1975, 3 ♀ (ES); Dulan-Khan, Ero-Gol River [106°30'E 49°56'N], 4.VIII 1975, 2 ♀, ♂ (MK); 20 km NNE Darkhan [105°60'E 49°25'N], 5.VIII 1975, ♀, 2 ♂ (ES); *Ulaanbaatar* [106°59'E 47°45'N], 14.VII 1967, ♂ (VZ); 1.VII 1967, ♂ (IK); *Tuv Aimag*: Kerulen [108°30'E 47°50'N], 15.VIII 1976, 3 ♂ (IK); 17.VIII 1976, 2 ♀, ♂ (MK); *Khentii Aimag*: 25 km NE Bayan-Adraga [111°50'E 48°30'N], 6.VII 1976, 9 ♂ (MK); 6-7.VII 1976, 3 ♂ (IK); *Dornod Aimag*: 32 km SE Salkhit Mt., Numregin-Gol River [119°45'E 47°00'N], 22-23.VII 1971, ♂ (MK); 20 km ENE Bayan-Ula [113°30'E 49°10'N], 8.VII 1976, ♂ (IK); 60 km ENE Bayan-Burd, Derkhin-Tsagan-Obo Mt. [119°00'E 46°80'N], 2.VIII 1976, 2 ♂ (MK); 10 km SE Salkhit Mt., Dege-Gol River [119°30'E 47°10'N], 5.VIII 1976, ♂ (MK); *Sukhbaatar Aimag*: Dzotol [115°10'E 45°90'N], 12.VII 1971, 3 ♂ (IK); Dzotol-Khan-Ula Mt. [114°00'E 45°33'N], 20-21.VII 1976, 4 ♀, 2 ♂ (MK); 65 km NNW Darigang [113°10'E 45°55'N], 14.VII 1976, 2 ♂ (MK); 9 km SW Darigang [114°05'E 45°33'N], 6.VII 1971, 2 ♂ (MK).

DISTRIBUTION. Mongolia (Usv Aimag, Khovd Aimag, Zavkhan Aimag, Govi-Altai Aimag, Khuvsgul Aimag, Arkhangai Aimag, Bayan-Khongor Aimag, Uvurkhangai Aimag, Selenge Aimag, Ulaanbaatar, Tuv Aimag, *Khentii Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (European part, Siberia, Far East), China (Beijing, Hebei, Inner Mongolia, Qinghai, Shandong, Sichuan, Xinjiang, Xizang, Yunnan), Kazakhstan, Uzbekistan, Kyrgyzstan, Nepal.

Colletes conradti species-group

Colletes wacki Kuhlmann, 2002

Colletes wacki Kuhlmann in Kuhlmann & Dorn, 2002: 102–103, ♂ (holotype: ♂, 30 km S Ulangom, Uvs Aimag, Mongolia); Kuhlmann, 2005: 1381; Kuhlmann & Proshchalykin, 2013a: 266; Proshchalykin & Kuhlmann, 2015: 335.

Colletes conradti: Kuhlmann & Dorn, 2002: 99; Kuhlmann, 2005: 1379; 2009a: 30; Kuhlmann & Proshchalykin, 2013a: 264.

SPECIMENS EXAMINED. **Mongolia.** *Umnugovi Aimag*: Tost-Ula, Tsagan-Ders-Khuduk [100°53'E 43°28'N], 22.VI 1973, ♂ (GM); *Dornod Aimag*: Khukh-Nur Lake [115°40'E 49°35'N], 25.VI 1976, ♀, 8 ♂ (MK).

DISTRIBUTION. Mongolia (Uvs Aimag, Zavkhan Aimag, Govi-Altai Aimag, Uvurkhangal Aimag, Umnugovi Aimag, Ulaanbaatar, Tuv Aimag, *Dornod Aimag), Russia (Eastern Siberia), China Hebei, Inner Mongolia, Qinghai), Kazakhstan.

Colletes marginatus species-group

Colletes chengtehensis Yasumatsu, 1935

Colletes chengtehensis Yasumatsu, 1935: 1–2, 35–36, ♀, ♂ (type locality: Jehol, Cheng-teh, China); Kuhlmann, 2009a: 30; 2009b: 18; Kuhlmann & Proshchalykin, 2011: 19; 2013a: 265; 2013b: 434; Proshchalykin & Kuhlmann, 2012: 25; 2015: 339; Niu *et al.*, 2014: 468.

Colletes pallescens: Kuhlmann & Dorn, 2002: 99; Kuhlmann, 2005: 1379.

SPECIMENS EXAMINED. **Mongolia.** *Usv Aimag*: 30 km NE Barun-Turun [94°24'E 49°39'N], 5.VII 1968, 2 ♀ (MK); *Khovd Aimag*: 5 km SW Altaj-som, Bodonchin-Gol River [92°25'E 45°81'N], 4.VIII 1968, ♀ (MK); 22.VII 1970, ♂ (MK); 20 km SE Altaj, Elkhon [92°45'E 45°80'N], 27.VII 1970, ♂ (VZ); 25 km N Bulgan, Ulyastain-Gol River [92°10'E 46°07'N], 31.VII 1970, 3 ♂ (MK); 20 km N Bulgan, Ulyastain-Gol River [92°10'E 46°05'N], 30.VI–1.VII 1980, 7 ♂ (IK); *Khuvsgul Aimag*: 25 km W Tevshrulekh [101°75'E 47°10'N], 11.VIII 1970, 2 ♂ (MK); 17 km N Shine-Ider [99°50'E 49°10'N], 22.VII 1975, ♀ (EN); 20 km SE Toson-Tsengel, Selenga River [100°50'E 49°25'N], 25.VII 1975, 2 ♀, 7 ♂ (MK); 25.VII 1975, ♂ (ES); 25.VII 1975, 5 ♂ (EN); *Bayan-Khongor Aimag*: Orog-Noor Lake [100°42'E 45°05'N], 15–16.VIII 1967, ♂ (VZ); Orog-Noor Lake, near Tuin-Gool River [100°42'E 45°03'N], 5.VIII 1969, ♂ (IK); 60 km SW Shine-Dzhinst, Burkhan-Bulak [98°62'E 44°45'N], 29.VIII 1970, ♂ (EN); *Uvurkhangai Aimag*: Tatsyn-Tsagan-Noor Lake [101°75'E 45°46'N], 2–4.VIII 1969, ♂ (MK); *Selenge Aimag*: Ero-Gol River [106°91'E 49°72'N], 26.VII 1970, ♂ (MK); 25 km E Darkhan, Sharyn-Gol Mt. [106°15'E 49°10'N], 30.VII 1975, 2 ♀ (ES); Dulan-Khan, Ero-Gol River [106°30'E 49°56'N], 4.VIII 1975, ♂ (MK); 4.VIII 1975, ♀, 2 ♂ (ES); 4.VIII 1975, ♂ (EN); *Khentii Aimag*: 12 km N Gal-Shara [110°85'E 46°71'N], 30.VII 1971, 1 ♀, 6 ♂ (MK); 25 km NE Bayan-Adraga [111°50'E 48°30'N], 6–7.VII 1976, 14 ♀,

5 ♂ (IK); *Dornogovi Aimag*: 10 km SW Sainshand, Tushilge Mt. [110°08'E 44°53'N], 1.VIII 1971, 8 ♂ (IK); 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, 6 ♂ (EN); *Dornod Aimag*: 60 km ENE Bayan-Burd, Derkhin-Tsagan-Obo Mt. [119°00'E 46°80'N], 21.VII 1971, 2 ♂ (MK); 2-4.VIII 1976, 5 ♀, 5 ♂ (MK); 75 km SW Salkhit, Ar-Bulak [118°30'E 46°86'N], 24.VII 1971, 2 ♂ (MK); 20 km NW Sangin-Dalai-Noor Lake [116°55'E 47°10'N], 25.VII 1971, m # (MK); E part of Buir-Nur Lake [117°40'E 47°47'N], 6-7.VI 1976, 5 ♂ (IK); 26-28.VII 1976, ♀ (IK); 15 km E Dzun-Bayan [110°12'E 44°26'N], 15.VIII 1975, 3 ♂ (ES); 10 km SE Salkhit Mt., Dege-Gol River [119°30'E 47°10'N], 5.VIII 1976, 6 ♂ (IK); 32 km SE Salkhit Mt., Numregin-Gol River [119°45'E 47°00'N] 23.VII 1971, 2 ♂ (MK); 25.VI 1976, ♂ (MK); 9.VIII 1976, ♂ (MK); *Sukhbaatar Aimag*: 20 km E Baruun-Urt [113°16'E 46°41'N], 14.VII 1971, ♂ (MK); 40 km SE Baruun-Urt [113°18'E 46°48'N], 14.VII 1971, ♂ (MK); 9 km SW Darigang [114°05'E 45°33'N], 8.VII 1971, 2 ♂ (MK); Dzotol-Khan-Ula Mt. [114°00'E 45°33'N], 20-21.VII 1976, 3 ♀, 7 ♂ (MK).

DISTRIBUTION. Mongolia (Usv Aimag, Khovd Aimag, Zavkhan Aimag, Govi-Altai Aimag, Khuvsgul Aimag, Arkhangai Aimag, Bayan-Khongor Aimag, Bulgan Aimag, Uvurkhangai Aimag, Umnugovi Aimag, Selenge Aimag, Tuv Aimag, Khentii Aimag, Dornogovi Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (European part, Siberia), China (Beijing, Hebei, Shanxi, Gansu, Inner Mongolia, Heilongjiang, Xinjiang), Kazakhstan, Uzbekistan, Kyrgyzstan, Turkmenistan, Tajikistan, Iran, Europe, Georgia, Azerbaijan.

Colletes succinctus species-group

Colletes arsenjevi Kuhlmann, 2006

Colletes arsenjevi Kuhlmann in Kuhlmann & Quest, 2006: 3, ♂ (type locality: Lazovsky Nature Reserve, Primorskiy Terr., Russia); Kuhlmann & Quest, 2006: 2; Proshchalykin, 2007: 878; 2012: 449; Kuhlmann, 2009a: 31; 2009b: 19; Kuhlmann & Proshchalykin, 2011: 20; 2013a: 265; 2013b: 436; Niu *et al.*, 2013: 178; 2014: 454; Proshchalykin & Kuhlmann, 2015: 340.

Colletes succinctus: Kuhlmann & Dorn, 2002: 100.

SPECIMENS EXAMINED. **Mongolia.** *Arkhangai Aimag*: Ugij-Noor Lake [102°46'E 47°45'N], 1.IX 1967, ♀ (VZ); 15 km W Teryat, Tsagan-Nur Lake [99°65'E 48°20'N], 19.VII 1980, ♂ (MK); *Khentii Aimag*: 8 km S Norovlin [111°59'E 48°40'N], 25.VIII 1975, ♀ (MK); *Dornogovi Aimag*: 40 km W Erdene [110°20'E 44°50'N], 14.VIII 1975, ♂ (MK); 70 km ENE Sain-Shand, Dolotyn-Khuduk [111°10'E 44°45'N], 17.VIII 1975, ♂ (EN); 10 km E Dzun-Bayan [110°03'E 44°18'N], 15.VIII 1975, ♂ (ES); 13 km W Dash-Balbar, Uldza River [114°00'E 49°20'N], 24.VIII 1975, 5 ♂ (EN); *Sukhbaatar Aimag*: 80 km NNE Barun-Urt [114°00'E 47°10'N], 19.VIII 1975, ♀ (ES).

DISTRIBUTION. Mongolia (Arkhangai Aimag, *Khentii Aimag, *Dornogovi Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (Far East), China (Shanxi, Gansu, Inner Mongolia, Xinjiang), Kazakhstan.

Colletes collaris Dours, 1872

Colletes collaris Dours, 1872: 296, ♀, ♂ (type locality: “Cannes, Algére, îles de l’Archipel grec”); Kuhlmann & Dorn, 2002: 100; Kuhlmann, 2005: 1379; 2009a: 31; Proshchalykin, 2007: 878; 2012: 449; Kuhlmann & Proshchalykin, 2011: 21; 2013a: 265; 2013b: 436; Niu *et al.*, 2013: 181; 2014: 478.

SPECIMENS EXAMINED. **Mongolia.** *Arkhangai Aimag:* Tsagan-Noor Lake, 18 km NE Dash-Baldooro [99°83'E 48°38'N], 23.VII 1975, 2 ♂ (MK); *Tuv Aimag:* Kerulen [108°30'E 47°50'N], 15.VIII 1976, ♂ (MK); *Dornogovi Aimag:* 70 km ENE Sain-Shand, Dolotyn-Khuduk [111°10'E 44°45'N], 17.VIII 1975, ♂ (EN); *Dornod Aimag:* 55 km NNE Khavirga [116°00'E 48°55'N], 21.VIII 1975, 2 ♂ (MK); 13 km W Dash-Balbar, Uldza River [114°00'E 49°20'N], 24.VIII 1975, ♀, ♂ (EN).

DISTRIBUTION. Mongolia (Arkhangai Aimag, Tuv Aimag, Khentii Aimag, *Dornogovi Aimag, Dornod Aimag, Sukhbaatar Aimag), Russia (forest zone of European part, Siberia, Far East), China (Beijing, Hebei, Jiangxi, Inner Mongolia, Heilongjiang, Qinghai, Xizang, Sichuan, Yunnan), Kazakhstan, South Korea, Japan (Honshu).

ACKNOWLEDGEMENTS

The work of first author was supported by President grant for government support of the Leading Scientific Schools of the Russian Federation (No HII-150.2014.4) and the grants of the Russian Funds for Basic Research (No 14-04-00649, No 15-29-02466 офи_м).

REFERENCES

- Astafurova, Yu.V. & Proshchalykin, M.Yu. 2015. New and little known bees of the genus *Sphecodes* Latreille (Hymenoptera: Halictidae) from Mongolia. *Far Eastern Entomologist*, 289: 1–9.
- Dours, J.A. 1872. Hyménoptères nouveaux du bassin méditerranéen. *Revue et Magasin de Zoologie*, 23: 293–311.
- Emeljanov, A.F., Kerzhner, I.M. & Kozlov, M.A. 1973. Joint Soviet-Mongolian investigations of the insect fauna of the Mongolian People's Republic during 1968 and 1971. *Entomologicheskoe Obozrenie*, 52(2): 466–482. [In Russian].
- Emeljanov, A.F., Kerzhner, I.M. & Kozlov, M.A. 1977. Joint Soviet-Mongolian investigations of the insect fauna of the Mongolian People's Republic during 1975 and 1976. In: Kerzhner, I.M. (Ed.). *Insects of Mongolia. Vol. 5.* Nauka, Leningrad. 7–16. [In Russian].
- Emeljanov, A.F. & Kozlov, M.A. 1978. Joint Soviet-Mongolian investigations of the insect fauna of the Mongolian People's Republic during 1978. In: Kerzhner, I.M. (Ed.). *Insects of Mongolia. Vol. 7.* Nauka, Leningrad. 7–13. [In Russian].
- Emeljanov, A.F., Zaytsev, V.F. & Kerzhner, I.M. 1968. Entomological expedition of the Zoological Institute, USSR Academy of Sciences, to the Mongolian People's Republic in 1967. *Entomological Review*, 47(4): 575–579.
- Eversmann, E. 1852. Fauna Hymenopterologica Volgo-Uralensis. *Bulletin de la Imperiale Society d'Naturalistes de Moscou*, 25(2): 3–137.

- Friese, H. 1913. Vorläufige Diagnosen von neuen Bienenarten, die von den Expeditionen Roborovsky-Koslov (1893–95) und von Koslov (1899–1901) aus Centralasian mitgebracht wurden und im Zoologischen Museum der Kaiserl. Akademie der Wissenschaften in St. Petersburg aufbewahrt werden. *Annuaire de Musée Zoologique de l'Academie Imperiale des Sciences de St. Peterbourg*, 18: 59–61.
- Insects of Mongolia. In 11 Volumes.* Leningrad: Nauka. Vol. 1. 1972: 1–990; Vol. 2. 1974: 1–439; Vol. 3. 1975: 1–675; Vol. 4. 1976: 1–639; Vol. 5. 1977: 1–757; Vol. 6. 1979: 1–512; Vol. 7. 1980: 1–576; Vol. 8. 1982: 1–576; Vol. 9. 1984: 1–576; Vol. 10. 1989: 1–623; Vol. 11. 1990: 1–624. [In Russian].
- Kerzhner, I.M., Kozlov, M.A. & Korotyaev, B.A. 1982. Joint Soviet-Mongolian investigations of the insect fauna of the Mongolian People's Republic during 1980 and 1981. In: Kerzhner, I.M. (Ed.). *Insects of Mongolia. Vol. 8.* Nauka, Leningrad. 7–15. [In Russian].
- Kuhlmann, M. 2000. Katalog der paläarktischen Arten der Bienengattung *Colletes* Latreille, mit Lectotypenfestlegungen, neuer Synonymie und der Beschreibung von zwei neuen Arten (Hymenoptera: Apidae: Colletinae). *Linzer biologische Beiträge*, 32(1): 155–193.
- Kuhlmann, M. 2002. Neue Arten der Bienengattung *Colletes* Latreille aus Südtibet mit Beschreibung der Route der „Deutschen Tibet Expedition Ernst Schäfer 1938/39“ (Hymenoptera: Apidae: Colletinae). *Linzer biologische Beiträge*, 34(2): 1155–1178.
- Kuhlmann, M. 2005. Faunistik und Zoogeographie der Bienengattung *Colletes* Latreille 1802 (Hymenoptera: Apidae: Colletinae) in Mittelasien. *Linzer biologische Beiträge*, 37(2): 1353–1396.
- Kuhlmann, M. 2009a. Erster Nachtrag zur Kenntnis der Bienengattung *Colletes* Latreille 1802 in der Mongolei mit Beschreibung einer neuen Art (Hymenoptera, Apiformes, Colletidae). *Beiträge zur Entomologie*, 59: 19–32.
- Kuhlmann, M. 2009b. Bees of the genus *Colletes* (Hymenoptera, Colletidae) from Central Asia collected by the Kyushu University expeditions. *Esakia*, 49: 15–20.
- Kuhlmann, M., Almeida, E.A.B., Laurenne, N. & Quicke, D.L.J. 2009. Molecular phylogeny and historical biogeography of the bee genus *Colletes* Latreille, 1802 (Hymenoptera: Apiformes: Colletidae), based on mitochondrial COI and nuclear 28 S sequence data. *Insect Systematics and Evolution*, 40: 291–318. DOI: 10.1163/139956009x12475840653733
- Kuhlmann, M. & Dorn, M. 2002. Die Bienengattung *Colletes* Latreille, 1802 in der Mongolei sowie Beschreibungen neuer Arten aus Sibirien und den Gebirgen Zentralasiens (Hymenoptera, Apidae, Colletinae). *Beitrag Entomologische Keltern*, 52(1): 85–109.
- Kuhlmann, M. & Proshchalykin, M.Yu. 2011. Bees of the genus *Colletes* Latreille 1802 of the Asian part of Russia, with a key to species (Hymenoptera, Apoidea: Colletidae). *Zootaxa*, 3068: 1–48.
- Kuhlmann, M. & Proshchalykin, M.Yu. 2013a. The bees of the genus *Colletes* Latreille 1802 of Mongolia (Hymenoptera, Apoidea: Colletidae). *Beiträge zur Entomologie*, 63(2): 255–269.
- Kuhlmann, M. & Proshchalykin, M.Yu. 2013b. The genus *Colletes* (Hymenoptera: Apoidea: Colletidae) in Central Asia. *Zootaxa*, 3750(5): 401–449. DOI: 10.11646/zootaxa.3750.5.1
- Kuhlmann, M. & Proshchalykin, M.Yu. 2014a. The bees of the genus *Colletes* Latreille 1802 of the European part of Russia, with keys to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, 3846(4): 591–596. DOI: 10.11646/zootaxa.3878.3.1
- Kuhlmann, M. & Proshchalykin, M.Yu. 2014b. New and little known bees of the genus *Colletes* Latreille 1802 (Hymenoptera: Colletidae) from Central Asia. *Zootaxa*, 3846(4): 591–596. DOI: 10.11646/zootaxa.3846.4.8
- Kuhlmann, M. & Quest, M. 2006. A new species of the bee genus *Colletes* Latreille, 1802 (Hymenoptera: Colletidae) from the Russian Far East and Mongolia. *Far Eastern Entomologist*, 157: 1–4.

- Morawitz, F. 1893. Supplement zur Bienenfauna Turkestans. *Horae Societatis Entomologicae Rossicae*, 28(1/2): 3–87.
- Niu, Z.-Q., Kuhlmann, M. & Zhu, C.-D. 2013. A review of the *Colletes succinctus*-group (Hymenoptera: Colletidae) from China with redescription of the male of *C. gigas*. *Zootaxa*, 3626(1): 173–187. DOI: 10.11646/zootaxa.3626.1.7
- Niu, Z.-Q., Zhu, C.-D. & Kuhlmann, M. 2014. The bees of the *Colletes* (Hymenoptera: Apoidea: Colletidae) from China. *Zootaxa*, 3856(4): 451–483. DOI: 10.11646/zootaxa.3856.4.1
- Noskiewicz, J. 1936. Die Palearktischen *Colletes*-Arten. *Prace Naukowe Wydawnictwo Towarzystwa Naukowego we Lwowie*, 3: 1–531.
- Noskiewicz, J. 1959. Weitere Beiträge zur Kenntnis der paläarktischen Arten der Gattung *Colletes* Latr. (Hym., Apidae). *Polskie pismo entomologiczne*, 29: 35–53.
- Nylander, W. 1852. Reviso synoptica apum borealium, comparatis speciebus Europae mediae. *Notiser ur Sällskapets pro Fauna et Flora Fennica Förhandlingar*, 2: 225–348.
- Osytsnjuk, A.Z. & Romankova, T.G. 1995. Family Colletidae. P. 480–489. In: Lehr, P.A. (Ed.). *Key to the insects of Russian Far East. Vol. 4. Pt 1.* Nauka, St. Petersburg. [In Russian].
- Pesenko, Yu.A. 1977. Bees of the genus *Nomioides* Schenck (Hymenoptera, Halictidae) of Mongolia. P. 583–585. In: Kerzhner, I.M. (Ed.). *Insects of Mongolia. Vol. 5.* Nauka, Leningrad. [In Russian].
- Pesenko, Yu.A. 1984a. Bees of the genus *Halictus* Latreille s. str. (Hymenoptera, Halictidae) of Mongolia and north-western China, with a review of publications on the Halictini of this region and a revision of the subgenus *Prohalictus* of the World fauna. P. 446–481. In: Korotyaev, B.A. (Ed.). *Insects of Mongolia. Vol. 9.* Nauka, Leningrad. [In Russian].
- Pesenko, Yu.A. 1984b. Bees of the tribe Nomioidini (Hymenoptera, Halictidae) in the Mongolian fauna: additions and corrections. P. 482–483. In: Korotyaev, B.A. (Ed.). *Insects of Mongolia. Vol. 9.* Nauka, Leningrad. [In Russian].
- Proshchalykin, M.Yu. 2007. Family Colletidae. In: Lelej, A.S. (Ed.). *Key to the insects of Russian Far East. Vol. 4. Pt 5.* Vladivostok: Dalnauka, 878–883. [In Russian].
- Proshchalykin, M.Yu. 2012. Family Colletidae. In: Lelej, A.S. (Ed.). *Annotated catalogue of the insects of Russian Far East. Vol. I. Hymenoptera.* Vladivostok: Dalnauka, 449–451. [In Russian].
- Proshchalykin, M.Yu. & Kuhlmann, M. 2012. The bees of the genus *Colletes* Latreille 1802 of the Ukraine, with a key to species (Hymenoptera: Apoidea: Colletidae). *Zootaxa*, 3488: 1–40.
- Proshchalykin, M.Yu. & Kuhlmann, M. 2015. Additional records of the genus *Colletes* Latreille (Hymenoptera: Apoidea: Colletidae) from Siberia, with a checklist of Russian species. *Zootaxa*, 3949(3): 323–344. DOI: 10.11646/zootaxa.3949.3.2
- Radoszkowski, O. 1891. Révision des armures copulatrices des mâles des genre *Colletes*. *Horae Societatis Entomologicae Rossicae*, 25(1/2): 249–260.
- Smith, F. 1846. Description of the British species of Bees comprised in the genera *Colletes* of Latreille and *Macropis* of Klug with observations on their economy. *Zoologist*, 4: 1274–1281.
- Smith, F. 1869. Descriptions of Hymenoptera from Japan. *Entomologist*, 4: 205–208.
- Yasumatsu, K. 1935. Insect of Jehol (VIII) – Order Hymenoptera (II), Superfamily Apoidea. *Report of the first scientific expedition to Manchoukuo under the leadership of Shigeyasu Tokunaga, June-October 1933. Section V, Division I, Part XII, Article 67:* 1–47.