

# Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by  
Far East Branch of the  
Russian Entomological Society  
and Laboratory of Entomology  
Institute of Biology and Pedology,  
Vladivostok

---

Number 51: 1-7

ISSN 1026-051X

December 1997

---

## A NEW SUBSPECIES OF THE *PSEUDOPANTHERA* *MACULARIA* L. (LEPIDOPTERA: GEOMETRIDAE, ENNOMINAE) FROM SOUTH SIBERIA

**E. A. Beljaev**

*Institute of Biology and Pedology, Vladivostok-22, 690022, Russia*

The geometrid moth *Pseudopanthera macularia cryptica* **ssp.n.** is described from South Siberia. New subspecies easily distinguished from nominative one in appearance and the female genitalia, but similar to that in the male genitalia.

KEY WORDS: Lepidoptera, Geometridae, new subspecies, South Siberia.

**Е.А. Беляев. Новый подвид *Pseudopanthera macularia* L. (Lepidoptera, Geometridae, Енноминае) из Южной Сибири // Дальневосточный энтомолог. 1997. N 51. С. 1-7.**

Описана пяденица *Pseudopanthera macularia cryptica* **ssp.n.** из Южной Сибири. Новый подвид легко отличим от номинативного по внешнему виду и гениталиям самки, но сходен с ним по строению гениталий самца.

*Биолого-почвенный институт, Дальневосточное отделение Российской Академии наук, Владивосток-22, 690022, Россия.*

### INTRODUCTION

There are 8 species of the genus *Pseudopanthera* Hübner, [1823] in Palaearctic region from West Europe to North India and West China (Prout, 1915; Wehrli, 1940). But most of them considerably differ from the type species

*Pseudopanthera macularia* (Linnaeus) by wing pattern and composition of this genus needs to be revised. *P. macularia* is widely distributed from Europe to North Mongolia and Transbaikalia. This species is polymorphic in the wing pattern, therefore numerous aberrations were described from Europe (see: Prout, 1915; Wehrli, 1940). After examination of the males and females of *P. macularia* from various localities I found a high variability of their genitalia too. But the specimens from South Siberia have a constant differences in appearance and in genitalia structure that allow to select South Siberian populations as a new subspecies. All records of *P. macularia* from South Siberia and adjacent regions (Staudinger, 1892; Staudinger & Rebel, 1901; Prout, 1915; Djakonov, 1926; Viidalepp, 1975, 1979; Korshunov & Viidalepp, 1982) should be assigned to *P. macularia cryptica* ssp. n. The description of new subspecies is given below. Holotype is deposited in the Zoological Institute (Sankt-Petersburg), paratypes are in the same collection and in the Institute of Biology and Pedology (Vladivostok).

***Pseudopanthera macularia cryptica* Beljaev, ssp. n.**


Figs 1-3, 6-8

*Venilia macularia* (Linnaeus, 1758): Staudinger, 1892: 375; Staudinger & Rebel, 1901: 331 (part.).

*Pseudopanthera macularia* (Linnaeus, 1758): Prout, 1915: 343 (part.); Djakonov, 1926: 58; Viidalepp, 1975: 478; 1979: 797 (part.); Korshunov & Viidalepp, 1982: 106.

TYPE MATERIAL. Holotype: ♂, Russia, Irkutskaya oblast', Bunbui, 28. V 1915 (Valdaev). Paratypes: Russia: 1 ♂, Republic Gornyi Altai, Altaiskii reserve, Teletskoe Lake, 1.VI 1939 (coll. unknown); 1 ♂, same loc., Yurgi river, 1010-1180 m, 19.VI 1939 (coll. unknown); 1 ♀, "Altai, Chodra, 900 m", 27.VI 1949 (G. Dulkeit); 1 ♂, Krasnoyarskii krai, "Stolby" reserve, Mana river, 28. VI 1958 (G. Dulkeit); 1 ♀, same loc., 8.VII 1965 (G. Dulkeit); 1 ♂, Krasnoyarskii krai, "Zapadnyi Sayan" 26.V 1962 (coll. unknown); 1 ♂, Irkutskaya oblast, Bunbui, 24. V 1916 (Valdaev); 1 ♀, same loc., 4. V 1915 (Valdaev); 1 ♀, Irkutskaya oblast, "Angara" [Angarsk], 9.VII 1938 (Nadetskaya); 1 ♂, south of Republic Sakha-Yakutia, Chulman river, Titovo, 13.VI 1950 (coll. unknown).


DESCRIPTION. MALE. Length of forewing 12-14 mm. Labial palpi short, almost the same as eye in length. Frons and vertex greyish-yellow. Antenna filiform. Thorax greyish-yellow. Forewing moderately broad, with pointed apex and prominent outer margin, reddish-yellow, with dark-brown band-like transverse lines and spots; antemedial transverse line wide, slightly arched; medial line consists of two separate spots: one on the costal margin near discal spot (usually fused with the latter) and another one between veins  $Cu_1$  and  $A_2$  (usually touched or fused with posterior part of postmedial transverse line); postmedial line angled outwardly between veins  $M_1$  and  $M_2$  or on vein  $M_2$ , interrupted between veins  $M_3$  and  $Cu_1$ ; submarginal transverse line



Figs 1-3. *Pseudopanthera macularia cryptica* ssp. n., holotype, male genitalia. 1 – lateral view, 2 – ventral view, 3 – aedeagus.

consists of three large separate spots on the costal margin, on veins  $M_3$  and  $Cu_2$  and near tornus; fringe dark-brown from apex to vein  $M_3$ , beyond  $M_3$  with separate two dark-brown spots. Hindwing coloured as forewing, but without antemedial and medial lines, both postmedial and submarginal lines consist of three separate spots. Base of wings dotted by dark-brown, discal spots on both wings large, dark-brown, almost equal in size. Legs yellow, hind tibia with two pairs of long spurs, without hair pencil.

**MALE GENITALIA** (Figs 1-3). Tegumen long, with moderately wide lateral sides, distinctly separable from vinculum. Uncus long, conic, pointed to apex, moderately arched. Socii small, membranous. Gnathos heavy sclerotized, V-shaped, its lateral arms curved caudally, medial plate with sagittal row of



Figs 4-8. *Pseudopanthera macularia*, genitalia. 4, 5 – *Pseudopanthera m. macularia*, male genitalia (Germany): 4 – lateral view, 5 – aedeagus. 6-8 – *Pseudopanthera m. cryptica* ssp. n., paratypes, female genitalia: 6 – total view (Bunbui), 7, 8 – corpus bursae (7 – “Stolby” reserve, 8 – “Angara”).

strong thorns variable in number. Valva narrow, long, moderately sclerotized, slightly conic, costal margin heavy sclerotized, with short thorn at apex, base of valva with small coremata laterally. Transtilla long, narrow, medially interrupted, sharply angled caudally at middle. Juxta folded in two, deeply invaginated into abdomen, with pair of very long thin processes reaching level of uncus base and bearing widely spaced thorns on its distal part. Cristae as flat slightly sclerotized plate with long hair-like setae. Anellus membranous, with pair long triangular lateral plates placed near transtilla and covered by rare long setae. Vinculum long, relatively thin, with broad triangular extensions latero-ventrally. Aedeagus long, thin, slightly S-shaped, with strong sclerotized triangular extension at apex, vesica without cornuti.

FEMALE. Very similar to male in size and colour, but with thinner antenna.

FEMALE GENITALIA (Figs 6-8). Ovipositor typical medium-size, papillae anales moderately sclerotized, with minute longitudinal plication. Posterior apophyses relatively long, approximately two times as long as anterior apophyses. Eight segment widely sclerotized dorsally, and with membranous transverse plication ventrally. Ductus bursae heavy sclerotized, wide, long, almost cylindrical, its posterior margin continued to the sclerotized band-like lateral plates touching the eighth tergite anteriorly. Corpus bursae membranous, various in size and shape, with irregularly plicated sclerotized ring in posterior part. Signum quadrangular in base, flattened to the apex, with pair of lateral spines well developed.

DISTRIBUTION. Russia: south of the Central and East Siberia; probably North Mongolia.

ETYMOLOGY. *Cryptica* (Greek) - concealed, covert. The new subspecies is named "*cryptica*" because of similarity in male genitalia with nominative form.

REMARKS. I consider subspecific range of the described form as tentative. Differences between *P. m. macularia* and *P. m. cryptica* in the female genitalia and in wing pattern correspond to the specific for the groups of close species in Ennominae. But I couldn't find clear differences between discussed forms in the male genitalia (except small difference in shape of vinculum). Taxonomic range of *P. m. cryptica* ssp. n. can be changed on specific after examination of the preimaginal stages or finding of the sympatric populations of the form under discussion and typical *P. macularia*.


### ***Pseudopanthera macularia macularia* (Linnaeus, 1758)**

(Figs 4, 5, 9-12)

*Phalaena macularia* Linnaeus, 1758, Syst. Nat (edn 10) 1: 521.

MATERIAL. 1 ♂, 1 ♀, West France, Rennes (R. Oberthür); 1 ♀, Russia, Kirovskaya oblast, Vyatka, 18.VI 1910 (coll. unknown); 9 ♂, 10 ♀, Russia, Bryanskaya oblast, Belye Berega, 22.V-13.VI 1969, 1971, 1972 (A. Kurentsov, Tseitgamel).

DISTRIBUTION. Europe, South Ural, Caucas, Transcaucasian region, Asia Minor; Tien-Shan (? ssp.).



Figs 9-12. *Pseudopanthera macularia macularia*, female genitalia. 9 – total view (Vyatka), 10-12 – corpus bursae (10 – Germany; 11, 12 – Belye Berega).

**Key to subspecies of *Pseudopanthera macularia***

- 1(2). Wings yellow; dark-brown transverse lines and spots moderately wide, with dusting of yellow scales; antemedial line usually interrupted; dots at base of wings small and dark-brown. Male genitalia: vinculum with rounded latero-ventral extensions and with more or less curved lateral arms (Figs 4, 5). Female genitalia: corpus bursae with inflation of posterior margin of sclerotized ring and with signum irregular spined and rounded (Figs 9-11).  
 ..... *P. macularia macularia* (L.)

- 2(1) Wings reddish-yellow; dark-brown transverse lines and spots generally wider, without dusting of yellow scales; antemedial line usually not interrupted; dots at base of wings larger and, black-brown. Male genitalia: vinculum with more or less angular latero-ventral extensions and with almost straight lateral arms (Figs 1-3). Female genitalia: corpus bursae without inflation of posterior sclerotized ring and with signum quadrangular basally, bearing pair of lateral strong developed spines (Figs 6-8). . . . . *P. macularia cryptica* ssp. n.

#### ACKNOWLEDGEMENTS

I wish to express my grateful thanks to Dr D. Stüning (Museum König, Bonn) and Dr A. L. Lvovsky (Zoological Institute, St.Petersburg) for loaning the valuable material.

#### REFERENCES

- Djakonov, A.M. 1926. Zur kenntnis der Geometriden Fauna des Minussinsk-Bezirks (Sibirien, Ienissej Gouv.). – Jahrbuch des Martjanov'schen Staatsmuseum in Minussinsk (Sibirien) 4 (1): 1-78. (In Russian and German).
- Korshunov, Ju.P. & Viidalepp, J.R. 1982. [Geometrid moths (Lepidoptera, Geometridae) of Khakassia. Report 2.] In: Poleznye i vrednye nesekomye Sibiri. Nauka Publ., Novosibirsk: 101-107. (In Russian).
- Prout, L.B. 1912-1916. Die spannerartigen Nachtfalter. In: Seitz, A. [Ed.]. Die Grossschmetterlinge der Erde. Stuttgart, 4: i-v, 1-479, pls. 1-25.
- Staudinger, O. 1892. Lepidopteren des Kentei-Gebirges. – Deut. Ent. Ztscht. Iris 5: 300-393.
- Staudinger, O. & Rebel, H. 1901. Catalog der Lepidopteren des palaearktischen Faunengebietes. Berlin: 411pp.
- Viidalepp, J. 1975. [On the fauna of geometrid mots (Lepidoptera, Geometridae) of the Mongolian People's Republic]. In: Nasekomye Mongolii. 3. Nauka Publ., Leningrad: 438-490. (In Russian).
- Viidalepp, J. 1979. [List of the geometrid moths of the fauna of USSR. IV.] – Entomologicheskoe Obozrenie 58 (4): 782-798. (In Russian).
- Wehrli, E. 1939-1954. Subfamilie Boarmiinae. In: Seitz, A. [Ed.] Die Grossschmetterlinge der Erde. Stuttgart, 4 (Supplement): 254-766, pls. 19-53.