

<https://doi.org/10.25221/fee.350.3>

<http://urn:lsid:zoobank.org:pub:2FBF48C9-ED26-4041-AA0B-6D00C1EA5C8F>

**FIRST RECORD OF THE GENUS *CORENNYS* BATES, 1884
(COLEOPTERA: CERAMBYCIDAE) FROM RUSSIA**

A. I. Miroshnikov

Sochi National Park, Moskovskaya str., 21, Sochi, 354002, Russia; Russian Entomological Society, Krasnodar, Russia. E-mail: miroshnikov-ai@yandex.ru

Summary. The genus *Corennys* Bates, 1884 is recorded from Russia for the first time based on the male of *C. sericata* Bates, 1884 collected in Kunashir Island, Russian Far East.

Key words: beetles, Cerambycidae, Lepturinae, *Corennys*, fauna, new record, Kunashir Island, Russian Far East.

А. И. Мирошников. Первая находка в России рода *Corennys* Bates, 1884 (Coleoptera: Cerambycidae) // Дальневосточный энтомолог. 2018. N 350. С. 17-19.

Резюме. Род *Corennys* Bates, 1884 впервые отмечается для фауны России по самцу *C. sericata* Bates, 1884 с острова Кунашир (Курильские острова).

The genus *Corennys* Bates, 1884 consists of 10 species distributed in Japan, China, Taiwan, northeastern India, Bhutan, and the countries of the Indochina Peninsula (Ohbayashi & Niisato, 2009; Löbl & Smetana, 2010). Until now the species of this genus has not been recorded from Russia. One specimen of *Corennys sericata* Bates, 1884 has been collected in Kunashir Island in 2015. Thus, both the genus *Corennys* and the species *C. sericata* are reported from Russia for the first time.

NEW RECORD

***Corennys sericata* Bates, 1884**

Fig. 1

Corennys sericata Bates, 1884: 225, pl. 1, fig. 2 (type locality: Japan: Nanai; Chiuzenji; Ontaki).

Corennys sericata sericata: Gressitt, 1951: 123; Wang & Hua, 2009: 167.

MATERIAL EXAMINED. **Russia:** Kunashir Island, southwestern shore of Goryachee Lake, 45°52'22"N / 145°29'11"E, 14–17.VII 2015, 1♂, leg. Yu. Sundukov & L. Sundukova (in coll. of A. Miroshnikov). **Japan:** Nagano, Kiriake, Akiyama, 20–22.VII 1996, 2♂, 2♀, leg. Shinji Nagai (in coll. of A. Miroshnikov ex coll. of N. Ohbayashi); Tochigi, Yumoto, 1.VIII 1960, 1♂, leg. H. Yokoyama (in coll. of A. Miroshnikov); Gunma Pref., Oku-Nikko, 8.VIII 1980, 2♂, 2♀, leg. K. Nagashima (in coll. of S. Murzin).

MEASUREMENTS. Male from Kunashir: body length 15 mm, humeral width 3.5 mm.

DISTRIBUTION. Russia: Kunashir Island (new record), Japan (Aurivillius, 1912; Winkler, 1929; Plavilstshikov, 1932; Matsushita, 1933; Mitono, 1940; Plavilstshikov, 1940; Kusama & Takakuwa, 1984; Hayashi & Villiers, 1985; Ohbayashi, 1992; Ohbayashi & Niisato, 2009).

NOTES. The records of *Corenmys sericata* from China (Gressitt, 1951; Hayashi & Villiers, 1985; Hua, 1982; 2002; Löbl & Smetana, 2010) and South Korea (Löbl & Smetana, 2010) require confirmation, whereas those from Laos (Hua, 1984) must be referred to another species.



Fig. 1. *Corenmys sericata* Bates, 1884, male from Kunashir Island (photo by K.V. Makarov).

ACKNOWLEDGEMENTS

I am grateful to Dr. K.V. Makarov (Moscow State Pedagogical University, Moscow, Russia) and Dr. Yu.N. Sundukov (State Nature Reserve “Kurilskiy”, Yuzhno-Kurilsk, Russia) who kindly provided the photo and specimen from Kunashir, to Dr. S.V. Murzin (Moscow, Russia) for the opportunity to study the specimens of *C. sericata* from his private collection, and to Dr. Nobuo Ohbayashi (Kamimiyada, Miura City, Japan) who generously presented me some specimens of *C. sericata*.

REFERENCES

- Aurivillius, C. 1912. Cerambycidae: Cerambycinae. P. 3–574. In: *Coleopterorum Catalogus, auspiciis et auxilio W. Junk, editus a S. Schenkling. Pars 39*. W. Junk, Berlin.
- Bates, H.W. 1884. Longicorn beetles of Japan. Additions, chiefly from the later collections of Mr. George Lewis; and notes on the synonymy, distribution, and habits of the previously known species. *The Journal of the Linnean Society of London. Zoology*, 18: 205–261 + pls 1–2.
- Gressitt, J.L. 1951. Longicorn beetles of China. In: Lepesme, P. (Ed.). *Longicornia. Études et notes sur les Longicornes. Vol. 2*. Paul Lechevalier, Paris. 667 pp. + 22 pls.
- Hua, L.-Z. 1982. *A check list of the longicorn beetles of China (Coleoptera: Cerambycidae)*. Zhongshan University, Guangzhou. 159 pp.

- Hua, L.-Z. 1984. *A list of the longicorn beetles of Laos (Coleoptera, Cerambycidae)*. Zhongshan University, Guangzhou. 155 pp.
- Hua, L.-Z. 2002. *List of Chinese Insects. 2*. Zhongshan (Sun Yat-sen) University, Guangzhou. 612 pp.
- Kusama, K. & Takakuwa, M. 1984. *The longicorn-beetles of Japan in color*. Japanese Society of Coleopterology, Kodansha (Tokyo). 565 pp. + 96 pls. [In Japanese]
- Löbl, I. & Smetana, A. 2010. *Catalogue of Palaearctic Coleoptera. Vol. 6. Chrysomeloidea*. Apollo Books, Stenstrup. 924 pp.
- Matsushita, M. 1933. Beitrag zur Kenntnis der Cerambyciden des japanischen Reichs. *Journal of the Faculty of Agriculture of the Hokkaido Imperial University*, 34(2): 157–445.
- Mitono, T. 1940. 94. Cerambycidae. In: Miwa, Y. & Chûjô, M. (Eds.). *Catalogus Coleopterorum Japonicorum. Pars 8*. Noda-Syobo, Taihoku. 283 pp.
- Ohbayashi, N., 1992. Taxonomic notes of Japanese Cerambycidae. *Acta Coleopterologica Japonica*, 2: 1–11.
- Ohbayashi, N. & Niisato, T. 2009. Review of the *Pyrocalymma* generic-group sensu Hayashi & Villiers, 1995, with descriptions of a new genus and two new species (Coleoptera, Cerambycidae, Lepturinae). *Special Bulletin of the Japanese Society of Coleopterology*, 7: 139–167.
- Plavilstshikov, N.N. 1932. Cerambycidae II. Teil. Cerambycinae: Cerambycini II [Hesperophanina, Phoracanthina, Ibdionina, Callidiopina, Graciliina, Obriina, Psebiina, Thraniina, Molorchina, Eroschemina, Pyrestina, Prothemina, Pytheina, Deilusina]. *Bestimmungstabellen der europäischen Coleopteren*, 102: 1–142.
- Plavilstshikov, N.N. 1940. *Longicorn beetles. Part 2. Fauna of the USSR. Insects, Coleoptera. Vol. 22*. USSR Academy of Sciences Press, Moscow & Leningrad. 785 pp. [In Russian]
- Winkler, A. 1929. Phytophaga. Cerambycidae. P. 1135–1226. In: *Catalogus Coleopterorum regionis palaearticae. 9–10*. Albert Winkler, Wien.