

SHORT COMMUNICATION

T. I. Arefina. *HYDROPTILA ITOI* KOBAYASHI, 1977, A NEWLY RECORDED CADDISFLY (TRICHOPTERA: HYDROPTILIDAE) FROM RUSSIA - Far Eastern Entomologist. 2002. N 112: 8.

Т. И. Арефина. *Hydroptila itoi* Kobayashi, 1977 – новый для России вид ручейников (Trichoptera: Hydroptilidae) // Дальневосточный энтомолог. 2002. N 112: 8.

A new for Russian fauna caddisfly species was found among material from Sakhalin Island provided by Dr. V. Labay (Laboratory of Applied Ecology of Sakhalin Research Institute of Fisheries and Oceanography, Yuzhno-Sakhalinsk, Russia). This species was described by Kobayashi from Hokkaido [1], later it has been recorded from Honshu [2].

Hydroptila itoi Kobayashi, 1977

MATERIAL: 5 pharate ♂, 27 larvae, Russia: south part of Sakhalin Island, Tunaicha Lake, 18.VII 2001 (V. Labay).

DISTRIBUTION. Russia (Sakhalin: new record), Japan (Hokkaido, Honshu).

1. Kobayashi, M. 1977. The list and new species of caddisflies from Hokkaido, Japan (Trichoptera, Insecta). – Bulletin of Kanagawa Prefecture Museum (Natural Sciences) 10: 1-14.

2. Ito, T., Tanida, K. & Nozaki, T. 1993. Checklist of Trichoptera in Japan. 1. Hydroptilidae and Lepidostomatidae. – The Japanese Journal of Limnology 54(2): 141-150.

Author's address:

Institute of Biology and Soil Science,
Vladivostok-22, 690022, Russia

© Far Eastern entomologist (Far East. entomol.) Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, Yu.A. Tshistjakov, N.V. Kurzenko

Address: Institute of Biology and Soil Science, Far East Branch of Russian Academy of Science, 690022, Vladivostok-22, Russia.

E-mail: entomol@ibss.dvo.ru

FAX: (4232) 310 193