

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 149: 1-4

ISSN 1026-051X

July 2005

NEW DATA ON KEROPLATIDAE (DIPTERA, SCIAROIDEA) FROM RUSSIAN FAR EAST

Alexander I. Zaitzev

*Moscow City Pedagogical University, Department of Biology, Faculty of Chemistry
and Biology, 111568, Moscow, Russia*

Monocentrota sidorenkoi sp. n. is described from South Primorye (Russia).
New data on morphology and distribution of *Platyura nigricoxa* (Okada) are given.

KEY WORDS: Diptera, Keroplatidae, *Monocentrota*, *Platyura*, new species,
Russian Far East.

**А. И. Зайцев. Новые данные по двукрылым семейства Keroplatidae
(Diptera, Sciaroidea) с Дальнего Востока России // Дальневосточный
энтомолог. 2005. N 149. С. 1-4.**

С юга Приморского края описан *Monocentrota sidorenkoi sp. n.* Приведены
новые данные по морфологии и распространению *Platyura nigricoxa* (Okada).

*Московский городской педагогический университет, химико-биологический
факультет, кафедра биологии, 111568, Москва, ул. Чечулина, 1, Россия.*

INTRODUCTION

Twenty seven species of Keroplatidae have been described previously from
Russian Far East (6 species of *Macrocerata* Meigen, 5 species of *Keroplatus* Bosc, 1
species of *Macrorrhyncha* Winnertz, 2 species of *Platyura* Meigen, 1 species of
Isoneuromyia Brunetti, 2 species of *Neoplatyura* Malloch, 2 species of *Urytalpa*
Edwards, 1 species of *Pyratula* Edwards, 7 species of *Orfelia* Costa (Zaitzev,
1999). In material, which was given to me by Dr. V.S. Sidorenko (Vladivostok)
who collected it in June 2001 in the south part of Primorskii krai, a new species of
genus *Monocentrota* Edwards was discovered, as well as one species, *Platyura*

nigricoxa (Okada), which was known earlier only by author's descriptions (Okada, 1937; 1938a, b; 1939).

All specimens including holotype are deposited in A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences (Moscow).

Present study have been supported partly by grant of Far Eastern Branch of the Russian Academy of Sciences «State value and dynamics of major components of the East Asian forest ecosystems» (S. Storozhenko, principal investigator).

DESCRIPTIONS OF SPECIES

Monocentrota sidorenkoi A. Zaitzev, sp. n.

Figs 1–3

MATERIAL. Holotype – ♂, Russia, Primorskii krai, Kamenushka, Light trap, 16.VII 2001 (Sidorenko leg.). Paratypes – 1 ♂, 3 ♀, the same label.

DESCRIPTION. MALE. Head brownish, clypeus yellow, mouth parts and palpi brown; antennae light brown, sixth flagellar segment 2 times as long as wide. Apical segment of flagellum 2 times as long as preapical segment. Thorax yellowish brown. Mesonotum yellow with brown lateral spots and three longitudinal stripes bearing bristles. Scutellum brownish with row of long bristles. Wings clear; wing length 3.0 mm; C extended distinctly more than halfway from R_5 to M_1 . Halteres yellow. Legs yellow; fore tibia 1.4 times as long as fore basitarsus; mid tibia 1.2 times as long as mid basitarsus; hind tibia 1.5 times as long as hind basitarsus. Abdomen brown; tergites with yellow fore and hind margins. Genitalia yellow.

FEMALE. Similar to male. Ovipositor light brown.

RELATIONSHIPS. This species is close to *M. lundstroemi* Edwards, from which it is distinguished by the structure of the genitalia.

ETYMOLOGY. New species is named in honour of Dr. V.S. Sidorenko (Vladivostok, Russia) who collected material.

Platyura nigricoxa (Okada, 1937)

Figs 4–6

Apemon similis Johnson f. *nigricoxa* Okada, 1937: 33.


Platyura nigricoxa: Krivosheina & Mamaev, 1988: 207.

MATERIAL. 5 ♂, Russia: Primorye, Kamenushka, Malaise trap, 30.VI 2001 (Sidorenko leg.).


MALE. Head black; clypeus protruded forward, narrowed toward apex; mouth parts yellow; two basal segments of palpi brown, other ones yellowish brown; second palpal with round sensillae on external surface; antennae dark brown, gradually narrowed toward apex; sixth flagellar segment as long as wide; apical flagellar segment 1.3 times as long as preapical segment. Thorax dark brown; mesonotum shining with short bristles on lateral parts with two rows short dorsocentral bristles; scutellum with short bristles. Wing length 6.0–6.5 mm; wing with slightly darkened apical part and with dark band between R_1 and M_1 or M_2 ; C ending at tip of R_5 ; basal

cell with weak longitudinal fold; all longitudinal veins with macrotrichia. Legs yellowish brown; coxa yellow with brown basal parts; fore and mid femora yellow, hind femora with dark spots; tibia and tarsi brownish; fore tibia with 1 spur; mid and hind tibia with 2 spurs; fore tibia 1.2–1.3 times as long as fore basitarsus; mid tibia 1.2 times as long as mid basitarsus; hind tibia 1.7 times as long as hind basitarsus. Abdomen dark brown; tergite IV with yellow spot; genitalia dark brown.

FEMALE unknown.



Figs 1–4. *Monocentrota sidorenkoi* sp. n. (1–3) and *Platyura nigricoxa* (Okada) (4): 1 – ovipositor; 2 – male genitalia ventrally; 3 – the same dorsally; 4 – wing.



Figs 5, 6. Male genitalia of *Platyura nigricoxa* Okada: 5 – dorsally; 6 – ventrally.

REMARKS. According Okada (1937; 1938a, b; 1939) *P. nigricoxa* is known from Hokkaido, South Kuril and Sakhalin. This species is distinguished from European *P. marginata* (Meigen) by the having of dark markings on the wing.

REFERENCES

- Krivosheina, N.P. & Mamaev, B.M. 1988. Family Keroplatidae. – In: Soós, A. & Papp, L. (eds.). Catalogue of Palaearctic Diptera. Vol. 3. Ceratopogonidae–Mycetophilidae. Budapest. Academiai Kiado: 199–210.
- Okada, I. 1937. Nachtrag zu den Nematoceren von dem Kurilen (Diptera). – Transactions of the Sapporo Natural History Society 15 (1): 33–39.
- Okada, I. 1938a. Einige Fungivoriden vom Dasetsu–Gebirge in Hokkaido. – Insecta Matsumurana 12 (2,3): 91–99.
- Okada, I. 1938b. Beitrag zur Kenntnis der Ceroplatinen–Fauna Japans (Dipt., Fungivoridae). – Insecta Matsumurana 13 (1): 17–32.
- Okada, I. 1939. Studien über die Pilzmücken (Fungivoridae) aus Hokkaido (Diptera, Nematocera). – Journal of Faculty of Agriculture Hokkaido University 49 (4): 267–336.
- Zaitzev, A.I. 1999. Fam. Keroplatidae. – Key to the insects of the Russian Far East. Vol. VI. Diptera and Siphonaptera. Pt. 1, Vladivostok, Dalnauka: 144–151.

© Far Eastern entomologist (Far East. entomol.) Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, V.S. Sidorenko, N.V. Kurzenko

Address: Institute of Biology and Soil Science, Far East Branch of Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: entomol@ibss.dvo.ru

FAX: (4232) 310 193