

### NEW HOST RECORDS FOR PARASITIC DIPTERANS OF THE GENUS *PHASIA* LATREILLE (DIPTERA: TACHINIDAE) FROM THE RUSSIAN FAR EAST

T. O. Markova

*Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far East Branch of the Russian Academy of Sciences, Vladivostok, 690022, Russia. E-mail: martania@mail.ru*

**Summary.** The present report provides new data on the hosts of *Phasia albopunctata* (Baranov), *Ph. aurigera* (Egger), *Ph. aurulans* Meigen, *Ph. barbifrons* (Girschner), *Ph. obesa* (F.), *Ph. hemiptera* (F.), *Ph. subcoleoptrata* (L.), *Ph. zimini* (D.-M.), and *Ph. pusilla* Meigen (Tachinidae: Phasiinae) from the Russian Far East.

**Key words:** Diptera, Phasiinae, Phasiini, host, Heteroptera, Primorsky Krai, Russia.

**Т. О. Маркова. Новые сведения о хозяевах паразитических двукрылых рода *Phasia* Latreille (Diptera: Tachinidae) на Дальнем Востоке России // Дальневосточный энтомолог. 2023. N 475. С. 31-36.**

**Резюме.** Приведены новые сведения о хозяевах *Phasia albopunctata* (Baranov), *Ph. aurigera* (Egger), *Ph. aurulans* Meigen, *Ph. barbifrons* (Girschner), *Ph. obesa* (Fabricius), *Ph. hemiptera* (F.), *Ph. subcoleoptrata* (Linnaeus), *Ph. zimini* (D.-M.) и *Phasia pusilla* Meigen (Tachinidae: Phasiinae) на Дальнем Востоке России.

#### INTRODUCTION

The *Phasia* Latreille, 1804, one of the largest genera in the subfamily Phasiinae, belongs to the tribe Phasiini. Members of this genus are distributed worldwide. The *Phasia* species parasitize Heteroptera including crop pests and predatory bugs (Dupuis, 1963; Draber-Mońko, 1965; Herting, 1971; Nishiyama *et al.*, 1995; Shima, 2006; Sun & Marschall, 2003; Aksenenko *et al.*, 2012; etc.). An updated list of the *Phasia* species from the Nearctic, Palearctic, Afrotropical, Australian, and Oriental realms and their associated hosts has been published elsewhere (Sun & Marschall, 2003).

In the Russian Far East fauna, the genus *Phasia* is represented by 11 species. The present report provides new and summarizes available data on the distribution and hosts of nine of them: *Phasia albopunctata* (Baranov, 1935), *Ph. aurigera* (Egger, 1830), *Ph. aurulans* Meigen, 1824, *Ph. barbifrons* (Girschner, 1887), *Ph. obesa* (Fabricius, 1798), *Ph. hemiptera* (Fabricius, 1794), *Ph. subcoleoptrata* (Linnaeus, 1767), *Ph. zimini* (Draber-Mońko, 1965), and *Phasia pusilla* Meigen, 1824. To date, there is still a lack of data on the hosts of *Phasia rohdendorfi* (Draber-Mońko, 1965). *Ph. takanoi* (Draber-Mońko, 1965) in Japan has been reared from *Menida scotti* Put. (Nishiyama *et al.*, 1995; Shima, 2006).

## MATERIAL AND METHODS

The information about the distribution of the species is according to the available publications (Ziegler & Shima, 1996; Draber-Mońko, 2012; Sun & Marschall, 2003; Ziegler & Shima, 2003; Richter, 2004, 2005; Cha & Han, 2009; O'Hara *et al.*, 2009; Zeegers, 2017). The techniques of bug collection and keeping, assessment of the infestation rate, and rearing of adult Phasiinae were described in detail earlier (Markova *et al.*, 2018). All the materials used are deposited in the author's private collection. The species are named according to the systematic lists published by the experts in the respective taxa (Herting, 1984; Thomas *et al.*, 1988; Aukema & Rieger, 1996, 1999, 2001; 2006; Vinokurov *et al.*, 2010). In the list below, *n* means number of specimens.

## RESULTS

### *Phasia albopunctata* (Baranov, 1935)

MATERIAL. Primorsky Krai: Ussuriysky Nature Reserve (43°40'00" N, 132°30'00" E), ex \**Carbula putoni* Jak. (Pentatomidae), August 25, 2001, 1 ♀ (T. Markova, M. Maslov).

DISTRIBUTION. Russia: Western and Eastern Siberia, Far East (Amur Oblast, Khabarovsk Krai, and Primorsky Krai); Japan (Hokkaido), Taiwan, Pakistan.

HOST. This species was reared ex *Dolycoris baccarum* (L.) (Pentatomidae) in Siberia and in Primorsky Krai (Kolomiets, 1976; Markova, 1999), ex *Palomena angulosa* (Motsch.), *P. viridissima* (Poda), *Picromerus bidens* (L.), *Pentatoma semiannulata* (Motsch.), *Lelia decempunctata* (Motsch.) (Pentatomidae) and *Coreus marginatus orientalis* (Kir.) (Coreidae) in Primorsky Krai (Markova, 1999; Markova *et al.*, 2020), and ex *Eysarcoris lewisi* Dist. (Pentatomidae) in Japan (Nishiyama *et al.*, 1995; Shima, 2006). *Carbula putoni* Jak. was first recorded as a host from Primorsky Krai. The duration of development in the puparium was 12 days (*n*=1).

### *Phasia aurigera* (Egger, 1830)

MATERIAL. Primorsky Krai: village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Palomena viridissima* (Poda) (Pentatomidae), August 17, 2017, 1 ♂ (T. Markova).

DISTRIBUTION. Russia: Far East (Khabarovsk Krai, Primorsky Krai); Western Europe, Korean Peninsula, China.

HOST. *Palomena viridissima* (Poda) was first recorded as a host from Primorsky Krai. The duration of development in the puparium was 10 days (*n*=1). In Western Europe and European Russia, this fly is known as a parasite of *Palomena prasina* (L.), *Rhaphigaster nebulosa* Poda (Pentatomidae) and *Coreus marginatus marginatus* (L.), *Gonocerus acuteangulatus* (Goeze), *G. juniperi* H.-S. (Coreidae) (Dupuis, 1963; Draber-Mońko, 1965; Herting, 1971; Kolomiets, 1976; Tschorsnig & Herting, 1994; San & Marshall, 2003).

### *Phasia aurulans* Meigen, 1824

MATERIAL. Primorsky Krai: village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Acanthosoma haemorrhoidalis angulatum* Jak. (Acanthosomatidae), August 17, 2018, 1 ♀ (T. Markova).

DISTRIBUTION. Russia: northern and northwestern European part, Siberia, Far East (Primorsky Krai, Amur Oblast); Western Europe, Kazakhstan, Japan (Hokkaido), USA, Canada.

HOST. *Acanthosoma haemorrhoidalis angulatum* Jak. was first recorded as a host from Primorsky Krai. The duration of development in the puparium was 11 days (*n*=1). In South Carolina (USA), this fly is known as a parasite of *Elasmucha lateralis* (Say) (Acanthosomatidae) (San & Marshall, 2003).

***Phasia barbifrons* (Girschner, 1887)**

MATERIAL. Primorsky Krai: village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Palomena viridissima* (Poda) (Pentatomidae), August 15, 2020, 1 ♀ (T. Markova).

DISTRIBUTION. Russia: middle of the European part, Western Siberia, Far East (Primorsky Krai); Western Europe, China, Vietnam.

HOST. *Palomena viridissima* (Poda) was first recorded as a host from Primorsky Krai. The duration of development in the puparium was 12 days ( $n=1$ ).

***Phasia hemiptera* (Fabricius, 1794)**

MATERIAL. Primorsky Krai: village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Acanthosoma haemorrhoidalis angulatum* Jak. (Acanthosomatidae), August 22, 2018, 1 ♀ (T. Markova).

DISTRIBUTION. Russia: northern and middle European part, Siberia, Far East (Amur Oblast, Khabarovsk Krai, Primorsky Krai, Kamchatka, Sakhalin Oblast); Western Europe, Transcaucasus; China, Korean Peninsula, Japan (Hokkaido, Honshu).

HOST. This species was reared ex *Pentatoma metallifera* (Motsch.) in Siberia and Primorsky Krai (Kolomiets, 1976; Draber-Moňko, 1965; Markova, 1999), ex *Dolycoris baccarum* (L.), *Palomena viridissima* (Poda) (Pentatomidae), *Elasmotherus brevis* Lindb. (Acanthosomatidae) (Markova, 1999; Markova *et al.*, 2017) in Primorsky Krai, ex *Palomena prasina* (L.), *Pentatoma rufipes* (L.), *P. metallifera* (Motsch.) in Western Europe (Dupuis, 1963; Draber-Moňko, 1965; Herting, 1971; San & Marshall, 2003), and ex *Pentatoma rufipes* (L.), *P. japonica* (Dist.), *Lelia decempunctata* (Motsch.), *Palomena angulosa* (Motsch.) (Pentatomidae) in Japan (Nishiyama *et al.*, 1995; Shima, 2006). *Acanthosoma haemorrhoidalis angulatum* Jak. was first recorded as a host from Primorsky Krai. The duration of development in the puparium was 14 days ( $n=1$ ).

***Phasia obesa* (Fabricius, 1798)**

MATERIAL. Primorsky Krai: the village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Eurygaster testudinaria* (Geoffr.) (Scutelleridae), August 21, 2019, 1 ♀ (T. Markova).

DISTRIBUTION. Russia: middle of the European part, Siberia, Far East (Amur Oblast, Khabarovsk Krai, Primorsky Krai); Western Europe, Middle East, Israel, Transcaucasus, Kazakhstan, Middle Asia, Mongolia, Japan (Hokkaido), North America.

HOST. This species was reared ex *Zicrona caerulea* (L.), *Neottiglossa pusilla* (Gmel.) (Pentatomidae), *Leptopterna dolabrata* (L.), *Lygus pratensis* (L.), *L. rugulipennis* Popp. (Miridae), *Myrmus miriformis* (Fall.) (Rhopalidae), and *Beosus maritimus* (Scop.) (Lygaeidae) in Western Europe (Dupuis, 1949, 1963; Draber-Moňko, 1965; Herting, 1971; San & Marshall, 2003). *Eurygaster testudinaria* (Geoffr.) was firstly recorded as a host from Primorsky Krai. The duration of development in the puparium was 15 days ( $n=1$ ).

***Phasia subcoleoptrata* (Linnaeus, 1767)**

MATERIAL. Primorsky Krai: village of Kaimanovka (43°37'49" N, 132°13'49" E), ex \**Dolycoris baccarum* (L.) (Pentatomidae), August 17, 2018, 1 ♀; ex \**Eurygaster testudinaria* (Geoffr.) (Scutelleridae), August 15, 2019, 1 ♀ (T. Markova).

DISTRIBUTION. Russia: southern and central European part, Eastern Siberia, Far East (Amur Oblast, Khabarovsk Krai, Primorsky Krai); Western Europe, Israel, Turkey, Iran, Transcaucasus, Kazakhstan, Middle Asia, North Africa.

HOST. In Western Europe and European Russia, this fly is known as a parasite of *Eurygaster integriceps* Put. (Scutelleridae), *Dolycoris baccarum* (L.), *D. numidicus* Horv., and *Stagonomus amoenus* (Brullé) (Pentatomidae) (Dupuis, 1963; Draber-Moňko, 1965; Herting, 1971; Kolomiets, 1976; Tschorsnig & Herting, 1994; San & Marshall, 2003). *Dolycoris baccarum* (L.) and *Eurygaster testudinaria* (Geoffr.) were first recorded as hosts from Primorsky Krai. The duration of development in the puparium ranged 12–13 days ( $n=2$ ).

#### ***Phasia zimini* (Draber-Moňko, 1965)**

MATERIAL. Primorsky Krai: Ussuriysky Nature Reserve (43°40'00" N, 132°30'00" E), ex \**Eysarcoris aeneus* (Scop.) (Pentatomidae), September 10, 2017, 1 ♂ (T. Markova, M. Maslov).

DISTRIBUTION. Russia: Far East (Amur Oblast, Khabarovsk Krai, Primorsky Krai).

HOST. This species was reared ex *Dolycoris baccarum* (L.) (Pentatomidae) in Primorsky Krai (Markova, 1999). *Eysarcoris aeneus* (Scop.) was firstly recorded as a host in Primorsky Krai. The duration of development in the puparium was 14 days ( $n=1$ ).

#### ***Phasia pusilla* Meigen, 1824**

MATERIAL. Primorsky Krai: Ussuriysky Nature Reserve (43°40'00" N, 132°30'00" E), ex \**Eysarcoris aeneus* (Scop.) (Pentatomidae), September 7, 2017, 1 ♂, September 8, 2017, 1 ♀ (T. Markova, M. Maslov).

DISTRIBUTION. Russia: northern and central European part, Siberia, Far East (Amur Oblast, Khabarovsk Krai, Primorsky Krai); Western Europe, Israel, Transcaucasus, Kazakhstan, Mongolia, Japan.

HOST. *Eysarcoris aeneus* (Scop.) was firstly recorded as a host from Primorsky Krai. The duration of development in the puparium ranged 10–11 days ( $n=2$ ). In Western Europe, European Russia, and Siberia, this fly is known as a parasite of *Chilacis typhae* (Perr.), *Nysius cymoides* (Spin.), *N. helveticus* (H.-S.) (= *N. lineatus* Costa), *Nithecus jacobaeae* (Schill.), *Cymus glandicolor* Hahn., *Kleidocerus ericae* (Horv.) (Lygaeidae); *Microporus nigrita* (F.) (Cydnidae); *Eysarcoris aeneus* (Scop.), *Eysarcoris ventralis* (Westw.) (= *Stollia inconspicua* (H.-S.)), *Bagrada hiliaris* (Burm.) (Pentatomidae); *Lyctocoris campestris* (F.), *Anthocoris sarothamni* Dg. & Sc., *A. nemoralis* (F.) (Anthocoridae), and *Prostemma aeneicolle* St. (Nabidae) (Viktorov & Kozharina, 1961; Dupuis, 1963; Draber-Moňko, 1965; Herting, 1971; San & Marshall, 2003; Aksenenko *et al.*, 2012).

#### **ACKNOWLEDGEMENTS**

I am grateful to E.V. Kanyukova (Vladivostok) for valuable advice during preparation of this paper, M.V. Maslov (Vladivostok) for his help in collecting material, and E.P. Shvetsov (Vladivostok) for his help in translating the paper into English. The research was carried out within the state assignment of Ministry of Science and Higher Education of the Russian Federation (theme No. 121031000151-3).

#### **REFERENCES**

- Aksenenko, E.V. Gaponov, S.P., Khitsova, L.N. & Kondratyeva, A.M. 2012. The predatory true bug *Prostemma aeneicolle* St. (Heteroptera, Nabidae) as a new host of the of tachinid fly *Phasia pusilla* Meig. (Diptera, Tachinidae, Phasiinae). *Bulletin of Moscow Society of Naturalist. Biological Series*, 117(5): 69–71. [In Russian]

- Aukema, B. & Rieger, C. (Eds.). 1996–2006. *Catalogue of the Heteroptera of the Palaearctic Region. Vol. 2–5*. Netherlands Entomological Society Amsterdam. Vol. 2, 1996, 361 pp.; Vol. 3, 1999, 577 pp.; Vol. 4, 2001, 346 pp.; Vol. 5, 2006, 550 pp.
- Cha, D.-J. & Han, H.-Y. 2009. Two species of the genus *Phasia* Latreille (Insecta: Diptera: Tachinidae) new to Korea. *Korean Journal of Systematic Zoology*, 25: 309–316.
- Dupuis, C. 1949. Contributions à l'étude des Phasiinae cimicophages. VII. Observations biologiques sur les parasites d' Hemipteres Hétéroptères à Richelieu (Indre-et-Loire) en 1946, 47, 48. *Annals de Parasitologie Humaine et Comparée*, 24: 211–242.
- Dupuis, C. 1963. Essai monographique sur les Phasiinae. *Mémoires du Muséum National d'Histoire Naturelle*, A(26): 1–461.
- Draber-Moňko, A. 1965. Monographie der paläarktischen Arten der Gattung *Alophora* R.-D. (Diptera, Larvaevoridae). *Annales Zoologici*, 23(6): 69–194.
- Draber-Moňko, A. 2012. State of knowledge of the tachinid fauna of Eastern Asia, with new data from North Korea. Part III. Phasiinae. Supplement. *Fragmenta Faunistica*, 55: 147–153.
- Herting, B. 1971. Arachnida to Heteroptera. P. 79–112. In: Herting, B. & Simmonds, F. (Eds.). *A Catalogue of parasites and predators of terrestrial arthropods. Section A. Host or prey/Enemy. Vol. I*. Farnham Royal, Commonwealth Agricultural Bureaux.
- Herting, B. 1984. Catalogue of Palaearctic Tachinidae (Diptera). *Stuttgarter Beiträge zur Naturkunde*, Serie A (Biologie), 369: 1–228.
- Kolomiets, N.G. 1976. Diptera of the subfamily Phasiinae (Tachinidae) in the fauna of Siberia and the Far East. P. 143–164. In: *Nasekomye Dalnego Vostoka*. DVNTs AN SSSR, Vladivostok. [In Russian]
- Markova, T.O. 1999. New host and distribution data of tachinid flies of subfamily Phasiinae (Diptera, Tachinidae) in Siberia and Russian Far East. *Far Eastern Entomologist*, 75: 1–8.
- Markova, T.O., Maslov, M.V. & Repsh, N.V. 2017. New data on biology and ecology of *Phasia hemiptera* (Diptera: Tachinidae, Phasiinae) in Primorskii krai. *A.I. Kurentsov's Annual Memorial Meetings*, 28: 95–100. [In Russian]
- Markova, T.O., Maslov, M.V. & Repsh, N.V. 2018. Modifications of rearing cages for insect research. *Euroasian Entomological Journal*, 17(5): 345–348. [In Russian] DOI: 10.15298/euroasentj.17.5.06
- Markova, T.O., Kanyukova, E.V. & Maslov, M.V. 2020. Ecology of the Dock Bug *Coreus marginatus orientalis* (Kir.) (Heteroptera, Coreidae), a Host of Parasitic Dipterans (Diptera, Tachinidae), in the South of Primorskii Territory. *Entomological Review*, 100(5): 620–628. DOI: 10.1134/S0013873820050048
- Nishiyama, M., Iwasa, M. & Hori, K. 1995. Parasitism by tachinid flies (Diptera, Tachinidae) of heteropterous insects in Tokachi, Hokkaido. *Japanese Journal of Entomology*, 63(1): 159–165.
- O'Hara J., Shima H. & Zhang, C. 2009. Annotated Catalogue of the Tachinidae (Insecta: Diptera) of China. *Zootaxa*, 2190: 1–236.
- Richter, V.A. 2004. 124. Sem. Tachinidae – Takhiny. P. 148–398. In: Lehr, P.A. (Ed.) *Key to the insects of Russian Far East, Vol. VI, Diptera and Siphonaptera, Part 3*. Dalnauka, Vladivostok. [In Russian].
- Richter, V.A. 2005. New data on the biology and distribution of Palaearctic Tachinidae (Diptera, Tachinidae). *Entomologicheskoe Obozrenie*, 84(4): 911–915. [In Russian, English translation *Entomological Review*, 85: 1227–1230].
- Shima, H. 2006. A host-parasite catalog of Tachinidae (Diptera) of Japan. *Makunagi (Acta Dipterologica)*, 2: 1–171.

- Sun, X. & Marschall, S. 2003. Systematics of *Phasia* Latreille (Diptera: Tachinidae). *Zootaxa*, 276: 320 pp.
- Thomas, J.H. & Richard, C. 1988. *Catalog of the Heteroptera, or True Bugs of Canada and the Continental United States*. Froeschner. Brill Academic Publishers.
- Tschorsnig, H. & Herting, B. 1994. Die Raupenfliegen (Diptera: Tachinidae) Mitteleuropas: Bestimmungstabellen und Angaben zur Verbreitung und Ökologie der einzelnen Arten. *Stuttgarter Beiträge zur Naturkunde*, Ser. A, 506: 1–170.
- Viktorov, G.A. & Kozharina, N.K. 1961. Trophic associations of some Phasiinae species (Diptera, Larvaevoridae) with the sunn pest and other bugs in Krasnodar Territory. *Zoologicheskii Zhurnal*, 40(1) 1: 52–59. [In Russian]
- Vinokurov, N.N., Kanyukova, E.V. & Golub, V.B. 2010. *Catalogue of the Heteroptera of Asian Part of Russia*. Nauka, Novosibirsk. 320 pp. [In Russian]
- Zeegers, T. 2017. Tachinid fauna (Diptera: Tachinidae) of Khabarovskii Krai, Russia. *Far Eastern Entomologist*, 330: 1–28.
- Ziegler, J. & Shima, H. 1996. Tachinid flies of the Ussuri area (Diptera: Tachinidae). *Beiträge zur Entomologie*, 46: 379–478.

---

© **Far Eastern entomologist (Far East. entomol.)** Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, S.A. Belokobylskij, M.G. Ponomarenko, V.A. Mutin, E.A. Beljaev, E.A. Makarchenko, A.V. Gorochoy, T.M. Tiunova, M.Yu. Proshchalykin, S.A. Shabalin, V.M. Loktionov

Address: Federal Scientific Center of the East Asia Terrestrial Biodiversity (former Institute of Biology and Soil Science), Far East Branch of the Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: storozhenko@biosoil.ru

web-site: <http://www.biosoil.ru/fee>