

Correspondence

<http://urn:lsid:zoobank.org:pub:E00F236A-7FF6-4C4B-8B1C-DDE0AFB4A556>

A. V. Volynkin^{1, 2 *}, M. S. Ivanova¹⁾. FIRST RECORD OF *XYLOMOIA RETINAX* MIKKOLA, 1998 (LEPIDOPTERA: NOCTUIDAE) FROM ALTAI REPUBLIC, RUSSIA. – *Far Eastern Entomologist*. 2016. N 324: 13-14.

1) *Altai State University, South Siberian Botanical Garden, Lenina pr. 61, Barnaul 656049, Russia.* *Corresponding author, E-mail: volynkin_a@mail.ru

2) *Tomsk State University, Laboratory of Biodiversity and Ecology, Lenina pr. 36, Tomsk 634050, Russia.*

Summary. *Xylomoia retinax* Mikkola, 1998 is recorded from the Russian Altai for the first time.

Key words: Lepidoptera, Noctuidae, Xyleninae, fauna, new record, Siberia, Russian Altai.

А. В. Вольткин, М. С. Иванова. Первое указание *Xylomoia retinax* Mikkola, 1998 (Lepidoptera: Noctuidae) для Республики Алтай, Россия // Дальневосточный энтомолог. 2016. N 324. С. 13-14.

Резюме. Впервые для фауны Российского Алтая приводится малоизвестный вид совок *Xylomoia retinax* Mikkola, 1998.

The fauna of noctuid moths (Lepidoptera: Noctuidae) of the Russian Altai was completely revised by Volynkin (2012) and included 583 species. Later, one more species was added by Volynkin (2015). During authors' field work in the northern part of Russian Altai in June 2016 one poorly-known species, *Xylomoia retinax*, is firstly found in the Altai Republic.

NEW RECORD

Family Noctuidae Latreille, 1809

Subfamily Xyleninae Guenée, 1837

Genus *Xylomoia* Staudinger, 1892

Xylomoia retinax Mikkola, 1998

Figs 1, 2


MATERIAL. Russia: Altai Republic, Shebalino district, SE vicinities of Kamlak vill., Gorno-Altai Botanical Garden, 51°37'30.49"N, 85°42'6.78"E, 400 m, meadow in mixed forest, 25–30.VII 2016, 2 ♂, coll. Volynkin A.V. & Ivanova M.S. (Deposited in the private collection of A.V. Volynkin, Barnaul, Russia).

DISTRIBUTION. European part of Russia, South Ural, West Siberia (Mikkola, 1998; Sviridov *et al.*, 2003; Pekarsky & Korb, 2012; Knyazev *et al.*, 2015).

NOTES. The species was described from two localities in West Siberia (Novosibirsk Region) and South Ural (Chelyabinsk Region) (Mikkola, 1998), and later was recorded from the European part of Russia (Yaroslavl' and Saratov regions) (Sviridov *et al.*, 2003; Pekarsky & Korb, 2012) and other localities in Omsk and Novosibirsk regions of West Siberia (Knyazev *et al.*, 2015). Here it is firstly recorded from Altai Republic.

ACKNOWLEDGEMENTS

We thank the Director of the Gorno-Altai Botanical Garden, Altanai Alekseevna Achimova, and the staff of the botanical garden for help with organization of the field research.



Figs 1–2. *Xylomoia retinax* from Altai Republic, Russia (with labels). 1 – adult male, dorsal view; 2 – male genitalia.

REFERENCES

- Knyazev, S.A., Ivonin, V.V., Dubatolov, V.V., Vasilenko, S.V. & Ponomaryov, K.B. 2015. New records of Lepidoptera from the South of West Siberia. *Amurian Zoological Journal*, 7(1): 43–50. [In Russian].
- Mikkola, K. 1998. Revision of the genus *Xylomoia* Staudinger (Lepidoptera: Noctuidae), with descriptions of two new species. *Systematic Entomology*, 23: 173–186.
- Pekarsky, O.N. & Korb, S.K. 2012. The first finding of *Xylomoia retinax* Mikkola, 1998 in Lover Volga region (Lepidoptera: Noctuidae). *Eversmannia*, 31–32: 115. [In Russian].
- Sviridov, A.V., Tikhomirov, A.M., Shutova, E.V., Blinushov, A.E., Kuznetsov, I.V., Bolshakov, L.V., Ryabov, S.A., Sitnikov, P.S. & Klepikov, M.A. 2003. Noctuid moths (Lepidoptera: Noctuidae) new for different areas of Russia, 1. *Russian Entomological Journal*, 11(4): 445–450. [In Russian].
- Volynkin, A. 2015. First record of *Nola crambiformis* Rebel, 1902 (Insecta: Lepidoptera, Nolidae) from the Russian Altai. *Check List*, 11(1): 1547.
- Volynkin, A.V. 2012. Noctuidae of the Russian Altai (Lepidoptera). *Proceedings of the Tigirek State Natural Reserve*, 5: 1–399.