

Correspondence

<http://urn:lsid:zoobank.org:pub:8A20FA19-C010-4766-961D-27A5127AB2D0>

V. S. Yakubovich^{1, *)}, E. S. Koshkin²⁾. **FIRST RECORD OF THE DRAGONFLY *AESCHNOPHLEBIA LONGISTIGMA* SELYS, 1883 (ODONATA: AESHNIDAE) FROM KHABAROVSKII KRAI. – Far Eastern Entomologist. 2016. N 324: 15-16.**

1) Far Eastern Medical University, Khabarovsk 680000, Russia. *Corresponding author E-mail: presid_11@mail.ru

2) Institute of Water and Ecological Problems, Far Eastern Branch of the Russian Academy of Sciences, Khabarovsk 680000, Russia. E-mail: ekos@inbox.ru

Summary. The dragonfly *Aeschnophlebia longistigma* Selys, 1883 is firstly recorded from Khabarovskii krai (Russia).

Key words: Odonata, Aeshnidae, dragonflies, fauna, new record, Khabarovskii krai, Russia.

В. С. Якубович. Е. С. Кошкин. Первая находка стрекозы *Aeschnophlebia longistigma* Selys, 1883 (Odonata: Aeshnidae) в Хабаровском крае // Дальневосточный энтомолог. 2016. N 324. С. 15-16.

Резюме. Стрекоза *Aeschnophlebia longistigma* Selys, 1883 впервые указывается из Хабаровского края (Россия).

INTRODUCTION

Nowadays the dragonfly *Aeschnophlebia longistigma* Selys, 1883 is known in Russia from the south part of Primorskii krai. This species was recorded as *A. kolthoffi* Sjösted, 1925 from Ussuriisk City (Belyshev, 1956), Putyatin Island (Belyshev, 1964), Barabashevka River (Belyshev, 1966), and Khanka Lake (Malikova & Ivanov, 2001). Endemic to Primorskii krai *A. zygoptera* Belyshev, 1965 was described from Tumannaya River near Khasan Lake (Belyshev, 1965). Now both *A. kolthoffi* and *A. zygoptera* are considered as synonyms of *A. longistigma* (Jiang, 2000; Malikova & Ivanov, 2001). We found *A. longistigma* in Khabarovsk City and it is the northernmost locality of this species in Russia.

NEW RECORD

Family Aeshnidae Rambur, 1842

Aeschnophlebia longistigma Selys, 1883

Fig. 1

MATERIAL EXAMINED. Russia: Khabarovskii krai, central part of Khabarovsk City, 11.VII 2016, 1 ♀ (E.S. Koshkin).

DISTRIBUTION. Russia: south part of Primorskii krai, Khabarovskii krai (new record); North and Central China, Korean Peninsula, Japan (Needham, 1930; Jiang, 2000; Lee, 2001; Malikova & Ivanov, 2001).

NOTES. The collected female is characterized by follow: length of hindwing 43 mm, abdomen length with appendages 49 mm; subcostal vein continues beyond the nodes only on forewing.



Fig. 1. *Aeschnophlebia longistigma* Selys, 1883, female from Khabarovsk, lateral view.

REFERENCES

- Belyshev, B.F. 1956. Materials to the Far East fauna of Odonata. *Trudy dalnevostochnogo filiala im. V.L. Komarova AN SSSR, seriya zoologicheskaya*, 3(6): 181–199. [In Russian].
- Belyshev, B.F. 1964. Contributions to the knowledge of the fauna of dragonflies (Odonata) of Siberia. I. *Fragmenta faunistica*, 11(5): 53–74. [In Russian].
- Belyshev, B.F. 1965. A new dragon-fly species (Odonata, Insecta) from the South Primorye. *Zoologicheskii Zhurnal*, 44(12): 1864–1866. [In Russian].
- Belyshev, B.F. 1966. Contributions to the knowledge of the fauna of dragonflies of Siberia. III. To the fauna and biology of dragonflies (Odonata) of Monkukhai rivulet from the South Primorye. *Fragmenta faunistica*, 11(5): 53–74. [In Russian].
- Jiang, Y.-H. 2000. Verification and Distribution of Genus *Aeschnophlebia* Selys (Odonata: Aeschnidae) in China. *Chinese Journal of Entomology*, 20: 63–67.
- Lee, S.-M. 2001. *The Dragonflies of Korean Peninsula (Odonata)*. Junghaeng-Sa, Seoul. 277 pp.
- Malikova, E.I. & Ivanov, P.Yu. 2001. The fauna of dragonflies (Insecta, Odonata) of Primorskii krai. P. 131–143. In: *Vladimir Ya. Levanidov's Biennial Memorial Meetings. Issue 1*. Dalnauka, Vladivostok. 376 pp. [In Russian].
- Needham, J.G. 1930. *A Manual of the Dragonflies of China: a monographic study of the Chinese Odonata*. The Fan Memorial Institute of Biology, Peiping. 345 pp.