

# **Far Eastern Entomologist**

Number 358: 24-28

ISSN 1026-051X

May 2018

<https://doi.org/10.25221/fee.358.3>

<http://urn:lsid:zoobank.org:pub:05B72E08-6D40-42CD-BA0B-965E74A82B6B>

## **FIRST RECORD OF THE FAMILY IBALIIDAE (HYMENOPTERA) FROM THE KURIL ARCHIPELAGO, RUSSIA**

**Yu. N. Sundukov<sup>1, 2)</sup>**

*1) Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far East Branch of the Russian Academy of Sciences, Vladivostok, 690022, Russia. E-mail: yun-sundukov@mail.ru*

*2) State Nature Reserve “Kurilskiy”, Yuzhno-Kurilsk, Sakhalinskaya oblast, 694500, Russia.*

**Summary.** Family Ibaliiidae, the genus *Ibalia* Latreille, 1802 and two species, *I. (Ibalia) leucospoides* (Hochenwarth, 1785) and *I. (Tremibalia) jakowlewi* Jacobson, 1899, are newly recorded from Kunashir Island (Kuril Archipelago). A key to the species of Ibaliiidae known from the Russian Far East is given also.

**Key words:** Hymenoptera, Ibaliiidae, fauna, new record, key, Kunashir Island, Russia.

**Ю. Н. Сундуков. Первая находка семейства Ibaliiidae (Нимфоптера) на Курильских островах, Россия // Дальневосточный энтомолог. 2018. N 358. С. 24-28.**

**Резюме.** Впервые с острова Кунашир (Курильские острова) указываются семейство Ibaliiidae, род *Ibalia* Latreille, 1802 и два вида, *I. (Ibalia) leucospoides* (Hochenwarth, 1785) и *I. (Tremibalia) jakowlewi* Jacobson, 1899. Приведена определительная таблица дальневосточных видов семейства Ibaliiidae.

## **INTRODUCTION**

Ibaliiidae is a small parasitoid family includes 20 recent species in three genera, *Ibalia* Latreille, 1802, *Heteribalia* Sakagami, 1949 and *Eileenella* Fergusson, 1992 in the World (Liu & Nordlander, 1994; Choi *et al.*, 2013; Aguiar *et al.*, 2013). All known Ibaliiidae are endoparasitoids of the wood wasp larvae (Hymenoptera: Siricidae) living in conifers and hardwoods (Liu & Nordlander, 1994; Hansen, 2010; Paretas-Martinèz *et al.*, 2013). Three species of Ibaliiidae are known from the Russian Far East, but this family has never been recorded from Kuril Archipelago (Melika, 2012). Present paper is based on material collected by author on Kunashir Island in 2013 and 2014.

## **NEW RECORDS**

### **Family Ibaliiidae**

#### **Genus *Ibalia* Latreille, 1802**

NOTES. There are three species of this genus in the fauna of the Russian Far East: worldwide *Ibalia (Ibalia) leucospoides* (Hochenwarth, 1785), Palaearctic *I. (Tremibalia) jakowlewi*

Jacobson, 1899, and *I. (T.) ornata* Belizin, 1968 known from Russia (Primorskiy kray) and China (Melika, 2012).

***Ibalia (Ibalia) leucospoides leucospoides (Hochenwarth, 1785)***

Fig. 1

*Ichneumon leucospoides* Hochenwarth, 1785: 345; type locality: not given in original description, but probably Berlin.

*Ichneumon cultellator* Fabricius, 1793: 167; type locality: “Germania”.

*Diplolepis ibalia* De Lamarck, 1817: 163; type locality: “France meridionale”.

*Ibalia suprunenkoi* Jacobson, 1899: 289; type locality: Russia, Sakhalin, between Douay harbor and Alexandrovsk.

*Ibalia arcuata* Dalla Torre & Kieffer, 1910: 23; type locality: “England, Italien, Deutschland”.

*Ibalia sachalinensis* Matsumura, 1911: 98; type locality: Russia, “Sakhalin, Mauka”.

*Ibalia picea* Matsumura, 1912: 163; type locality: Japan.

MATERIAL. **Russia:** South Kurile, Kunashir Island, Tretyakovo, dry fir, 21–22.VIII 2013, 2♀, leg. Yu. Sundukov and L. Sundukova.



Fig. 1. *Ibalia (Ibalia) leucospoides leucospoides*, female from Tretyakovo, Kunashir Island, lateral view.

DISTRIBUTION. Russia: Kunashir Island (new record), Sakhalin, Primorskiy kray; Japan, South Korea, China, Greece, Bulgaria, Hungary, Czech Republic, Poland, Sweden, Switzerland, Germany, Italy, Belgium, Netherlands, France, British Islands, Spain, Portugal, Turkey, Cyprus, Algeria, Tunisia, Morocco; introduced to Argentina, Australia, Tasmania, New Zealand (Liu & Nordlander, 1994; Muruaga de L'Argentier *et al.*, 2005; Ljubomirov *et al.*, 2006; Peeters, 2011; Melika, 2012; Choi *et al.*, 2013; Paretas-Martínez *et al.*, 2013).

HOSTS. Parasitoid of wood wasps of the genera *Sirex*, *Urocerus* and *Xeris* in coniferous forests (Liu & Nordlander, 1994; Choi *et al.*, 2013).



Fig. 2. *Ibalia (Tremibalia) jakowlewi*, female from Severyanka River, Kunashir Island, lateral view.

***Ibalia (Tremibalia) jakowlewi* Jacobson, 1899**

Fig. 2

*Ibalia jakowlewi* Jacobson, 1899: 288; type locality: “Iretj”, Balagansky district, Irkutskaya oblast, Russia.

*Ibalia takachihoi* Yasumatsu, 1937: 13; type locality: Mount Hikosan, Buzen, Kyushu, Japan.

MATERIAL. **Russia:** South Kurile, Kunashir Island, downstream Severyanka River, 5.VI 2014, 1♀, leg. Yu. Sundukov.

DISTRIBUTION. Russia: Kunashir Island (new record), Primorskiy kray, Irkutskaya oblast, Altai, European part; Japan, Korea, Georgia, Ukraine, Czech Republic, Slovakia, Poland, Austria, Italy, Germany, Finland (Liu & Nordlander, 1994; Martikainen & Viitasaari, 1996; Melika, 2012; Choi *et al.*, 2013; Dal Pos, 2017).

HOSTS. Parasitoid of wood wasps (Siricidae: Tremicinae) *Tremex longicollis* Konow, 1896 in Japan and *T. fuscicornis* (Fabricius, 1787) in Europe in hardwoods (Liu & Nordlander, 1994; Holý *et al.*, 2012; Choi *et al.*, 2013).

## KEY TO THE RUSSIAN FAR EAST SPECIES OF THE FAMILY IBALIIDAE

1. Eye length 2.3 times length of malar space. Male antenna usually with 13 flagellomeres. Dorsal pronotal area short. Propleuron weakly projecting ventrally. Anterior lateral crest of metacoxa triangular. Parasitoids of Siricinae in coniferous trees .....  
..... *Ibalia (Ibalia) leucospoides*
- Maximum length of eye more than 2.7 times length of malar space. Male antenna with 12 flagellomeres. Dorsal pronotal area large. Propleuron strongly projecting ventrally. Anterior lateral crest of metacoxa rounded. Parasitoids of Tremicinae in hardwood trees (Subgenus *Tremibalia*) ..... 2
2. Head and pronotum entirely black. Entire gena costate. Lateral sides of pronotum mainly rugose. Fore wing with narrower dark band behind submarginal cell .....  
..... *Ibalia (Tremibalia) jakowlewi*
- Head and pronotum at least partially yellow or brown. Upper part of gena smooth. Lateral sides of pronotum costate and often partially smooth. Fore wing with broad, indistinct dark area behind submarginal cell ..... *Ibalia (Tremibalia) ornata*

## REFERENCES

- Aguiar, A.P., Deans, A.R., Engel, M.S., Forshage, M., Huber, J.T., Jennings, J.T., Jonson, N.F., Lelej, A.S., Longino, J.T., Lohrmann, V., Mikó, I., Ohl, M., Rasmussen, C., Taeger, A. & Yu, D.S.K. 2012. Order Hymenoptera. In: Zhang Z.-Q. (Ed.). Animal Biodiversity: An outline of higher-level classification and survey of taxonomic richness (Addenda 2013). *Zootaxa*, 3703(1): 51–62.
- Choi, W.-Y., Lee, J.-W. & Suh, K.-I. 2013. Taxonomic review of the family Ibaliiidae (Cynipoidea: Hymenoptera) from Korea. *Entomological Research*, 43: 135–141.
- Dal Pos, D. 2017. First record of *Ibalia jakowlewi* Jacobson, 1899 and of the subgenus *Tremibalia* for Italy (Hymenoptera, Cynipoidea, Ibaliiidae). *Bollettino del Museo di Storia Naturale di Venezia*, 68: 53–57.
- Hansen, O.L. 2010. The family Ibaliiidae (Hymenoptera, Cynipoidea) in Norway. *Norwegian Journal of Entomology*, 57: 139–141.
- Holý, K., Psota, V., Šťastná, P. & Macek, J. 2012. Faunistic records from the Czech Republic and Slovakia (Hymenoptera: Ibaliiidae, Ichneumonidae). *Acta Musei Moraviae. Scientiae biologicae* (Brno), 97(2): 65–68.
- Liu, Z. & Nordlander, G. 1994. Review of the family Ibaliiidae (Hymenoptera: Cynipoidea) with keys to genera and species of the World. *Entomologica Scandinavica*, 25(4): 377–392.
- Ljubomirov, T., Raikova, M. & Georgiev, G. 2006. Ibaliiidae and Embolemidae (Insecta, Hymenoptera), new families to the fauna of Bulgaria. *Acta zoologica bulgarica*, 58(3): 425–430.
- Martikainen, P. & Viitasaari, M. 1996. The family Ibaliiidae (Hymenoptera, Cynipoidea) in Finland with *Ibalia jakowlewi* Jacobson new to Fennoscandia. *Sahlbergia*, 3: 24–27.
- Melika, Zh. 2012. Superfamily Cynipoidea. P. 139–146. In: Lelej A.S. (Ed.). *Annotated catalogue of the insects of Russian Far East. Volume I. Hymenoptera*. Dalnauka, Vladivostok. [In Russian]
- Muruaga de L'Argentier, S., Quintana de Quinteros, S., Gallardo, C. & Vilte, H. 2005. Establecimiento de *Ibalia leucospoides* (Hymenoptera: Ibaliiidae) endoparásito de *Sirex noctilio* (Hymenoptera: Siricidae) en La Almona (Jujuy, Argentina). *Revista de la Sociedad Entomológica Argentina*, 64(1–2): 23–25.

- Parets-Martinèz, J., Forshage, M., Buffington, M., Fisher, N., La Salle, J. & Pujade-Villar, J. 2013. Overview of Australian Cynipoidea (Hymenoptera). *Australian Journal of Entomology*, 52: 73–86.

Peeters, Th.M.J. 2011. *Ibalia leucospoides* in Nederland. *Hymeno Varia*, 3: 54–56.