



Original Article

Exploring fauna of Microlepidoptera in South Siberia: novel regional records and interception of quarantine species

Evgeny N. Akulov^a, Margarita G. Ponomarenko^{b,c,*}, Natalia I. Kirichenko^{d,e}^a All-Russian Plant Quarantine Center, Krasnoyarsk branch, Krasnoyarsk, 660075, Russia^b Federal Scientific Center of the East Asia Terrestrial Biodiversity FEB RAS, Vladivostok, 690022, Russia^c Far Eastern Federal University, Vladivostok, 690922, Russia^d Sukachev Institute of Forest SB RAS, Federal Research Center «Krasnoyarsk Science Center SB RAS», Krasnoyarsk, 660036, Russia^e Siberian Federal University, Krasnoyarsk, 660041, Russia

ARTICLE INFO

Article history:

Received 24 September 2019

Received in revised form

27 September 2019

Accepted 1 October 2019

Available online 9 October 2019

Keywords:

Microlepidoptera

New regional records

Quarantine species

Russia

Southern Siberia

ABSTRACT

We report the results of a faunistic study of Microlepidoptera performed in Southern Siberia using diverse sampling techniques allowing to cover various taxonomic groups. The provided taxonomic list is comprised of 64 species from 44 genera and 18 families, where all species represent novel geographical records. Of them, only 62 species inhabit Southern Siberia, the other two species, *Tuta absoluta* (Meyrick) and *Carposina sasakii* Matsumura are quarantine pests intercepted in this region. Eighteen genera and 3 families (Glyphipterigidae, Chimabachidae, and Oecophoridae) are reported for the first time from the studied region. Overall, 11 species from 7 families are novel for Siberia. Two species, *Bucculatrix panonica* and *Coleophora curictae*, are newly recorded in Russia. For the species newly recorded in the fauna of Siberia and Russia, male or female genitalia are illustrated. For *Coleophora curictae*, representing a novel record for Russia, the bionomics is given for the first time.

© 2019 National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA), Publishing Services by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Regular faunistic inventories of species essentially contribute to the knowledge of the regional fauna (Ponomarenko 2008, 2016; Akulov et al 2018; Knyazev et al 2018; Kirichenko et al 2019c) and allow to detect the species expanding their ranges, as well as intercept invasive and quarantine pests (Kenis et al 2007; Fattorini 2013; Akulov et al 2013, 2015).

Situated in the middle of Siberia, Krasnoyarskii Krai (its southern part) and the Republic of Khakassia, together often referred in Russian literature to South Siberia, represent an important transition zone for the species of organisms having their distribution on the west or on the east of the Palearctic zone. Lepidoptera is a highly diverse worldwide distributed group of insects (Kristensen et al 2007), among which a number of pests and invasive species

are also known (Suckling et al 2017). The faunal studies of this insect order have been conducted on the territory of South Siberia for already more than one century (see an extended literature review in Akulov et al 2018). However, these studies were sporadic, covering certain groups of Lepidoptera (mainly Macrolepidoptera), or were dedicated to biology and ecology of particular species having the pest status, which undoubtedly have an important value. For decades, Microlepidoptera have been paid much less attention.

Since 2000, we have been running faunistic studies on the south of Krasnoyarskii Krai and occasionally in the Republic of Khakassia gaining knowledge about the diversity of Microlepidoptera in the region, describing new species to science and developing regional DNA barcode libraries for certain microlepidopteran groups (Akulov et al 2013, 2014, 2015, 2018; Akulov and Kirichenko 2014; Kirichenko et al 2016, 2017a, 2017b, 2018, 2019a, 2019b, 2019c).

Here, we present the result of our most recent faunistic surveys of Microlepidoptera in South Siberia that were based on utilization of various sampling techniques, allowing to reveal the micromoths of different taxonomic groups, to clarify the species distribution at a regional and within-country scale, and to intercept quarantine pests.

* Corresponding author. Tel.: +7(904) 626 12 13; fax: +7(423)2 310 193.

E-mail addresses: akulich80@ya.ru (E.N. Akulov), margp@biosoil.ru (M. G. Ponomarenko), nkirichenko@yahoo.com (N.I. Kirichenko).

Peer review under responsibility of National Science Museum of Korea (NSMK) and Korea National Arboretum (KNA).

Material and methods

The species of Microlepidoptera were collected in the south of Krasnoyarskii Krai (15 localities) and in the Republic of Khakassia (2 localities) from April to August 2015–2019 (Figures 1 and 2). Overall, about 120 specimens of adult moths were collected. As an exception, 5 specimens of *Coleophora* spp. were sampled as larvae in the cases.

In Krasnoyarskii Krai, sampling was done in gardens (V.M. Krutovsky Botanical Garden, private gardens' community «KZK-2», private gardens' community «Pobeda», Opytnoe pole), city plantations (Krasnoyarsk), in green house complex and market with import fruit production, as well as in nature biotopes, particularly in the Nature reserve «Stolby», the Biryusa Bay, Pogorelskiy bor, and in the meadows and forest surrounding the settlements of Berezovskii, Novaya Syda, Kubekovo, and Minino. In the Republic of Khakassia, Microlepidoptera were collected in the Tuimskoe forestry and in the steppe (field research station of SIF SB RAS, Chjerno Ozero village) (Figures 1 and 2).

We used various sampling techniques in this study. Among them, the pheromone trapping was the most utilized method (Figure 2C). We used pheromone traps with pheromone for regional pests or quarantine species [*Grapholita molesta* (Busck, 1916), *Dendrolimus sibiricus* Tschetverikov, 1877, *Frankliniella occidentalis* (Pergande, 1895), *Lymantria dispar* (Linnaeus, 1758),

Carposina "niponensis" [niponensis auct. (nec Walsingham, 1900)], and *Ceratitis capitata* (Wiedemann, 1824)]. In pheromone traps, captured insects were fixed on the sticky substrate, which was replaced with a fresh one 1–2 times a week. The methodology for collecting material in pheromone traps has been described in detail in our recent study (Akulov et al 2018). We also used light trapping (the lamp DRV, 250W) (Figure 2F and H), sweeping moths from grass using the insect net, as well as making opportunistic sampling of moths resting on different surfaces. In addition, the two species of Coleophoridae were sampled as feeding larvae on leaves on *Ulmus pumila* and *Rosa* sp.

The moths from pheromone traps often had shabby wings because of sticky substrate in the traps. For their preservation, 70% ethanol was commonly used. The adult moths sampled by other approaches were all mounted and placed in dry collections. Coleophoridae larvae were preserved in 96% ethanol solution for DNA barcoding.

Identification of Microlepidoptera was done, where possible, based on external adult morphology and characters of their genital apparatus. Genitalia dissections and slide mounts were done following the study by Robinson (1976). Different keys (Lerh 1997, 1999; Medvedev 1981) and lepidopterological sites (Savela 2001; Lepiforum 2019; Moth Dissection UK Lepidoptera 2019; North American Moth Photographers Group 2019; Svenska fjärilar Lepidoptera 2019) were used for species determination. The two




Figure 1. Sampling locations in South Siberia. In the left corner on the map of Russia, the sampled area is shown by red square. Locations in Krasnoyarskii Krai: 1, Emeljanovskii District, the Biryusa Bay; 2, ibidem, Minino settl.; 3, Pogorelskiy bor, Arboretum SIF SB RAS; 4, Krasnoyarsk, private gardens' community «KZK-2»; 5, ibidem, private gardens' community «Pobeda»; 6, ibidem, Red Square; 7, Emeljanovskii District, Kubekovo settl.; 8, Krasnoyarsk, city plantations; 9, ibidem, market with imported fruits; 10, ibidem, greenhouse complex; 11, ibidem, Nature reserve «Stolby»; 12, ibidem, V.M. Krutovsky Botanical Garden; 13, Berezovskii District, Berezovskii settl.; 14, Krasnoturanskii District, Novaya Syda settl.; 15, Minusinskii District, Opytnoe pole settl.; locations in Khakassia Republic: 16, Shirinskii District, Tuimskoe forestry; 17, Shirinskii District, Chjerno Ozero vil., field research station of SIF SB RAS.




Figure 2. Some biotopes and sampling methods in South Siberia. In Krasnoyarskii Krai: A–B, V.M. Krutovsky Botanical Garden; C, pheromone trapping in private gardens' community «Pobeda»; D, Nature reserve «Stolby»; E, forests around Berezovskii settl.; F, light trapping in Minino settl. In Khakassia Republic: G, steppe around Chjernoje Ozero vil.; H, light trapping at the field research station of SIF SB RAS.

species of Coleophoridae, collected as larvae, were DNA barcoded in Forest zoology laboratory at Institut National de la Recherche Agronomique (INRA), Orleans (France), within postdoctoral fellowship of NIK. The Folmer fragment of the COI mtDNA gene, 658 bp, was targeted, using standard primers LCO (5' GGT CAA CAA ATC ATA AAG ATA TTG G 3') and HCO (5' TAA ACT TCA GGG TGA CCA AAA AAT CA 3') (Folmer et al 1994). DNA extraction, amplification, and sequencing conditions are given in the study by Kirichenko et al (2016). The DNA barcodes of *Coleophora curictaes* and *C. gryhipennella* and data on their vouchers are publicly available in the BOLD data set "DS- COLEOF", accessed at: dx.doi.org/10.5883/DS-COLEOF.

The taxonomic list is comprised exceptionally of the species that represented novel geographic records. The species that had been known from the region and that were recaptured in our surveys were not included in this article.

The map showing sampling localities in South Siberia is built using ArcGIS 10.6 (Esri, California, United States). The photographs of insects and genitalia were taken using a digital camera, Canon 650D (Japan) portable with the microscope Olympus CX21 (Japan).

For each species, details on the place and the date of collection, the number of specimens, and the gender are provided. The data on the trophic associations are compiled from the numerous literature and electronic resources cited in the essay for each species. Geographic range is indicated according to the area typification proposed by Gorodkov (1984, 1992). As the majority of listed species inhabit the temperate zone of the Old and New Worlds, we largely avoided repeating it every time when characterizing the species range. For each species, the detailed distribution in Russia is provided, except for quarantine species for which the whole actual range is listed. The names of the regions in Russia are given




Figure 3. Genitalia, ventral view: A, *Bohemannia pulverosella* (Stainton, 1849), female genitalia; B, *Monopis obviella* ([Denis et Schiffermüller], 1775), male genitalia; C, *Bucculatrix nigricomella* (Zeller, 1839); D, *Bucculatrix pannonica* Deschka, 1982, male genitalia; E–F, *Glyphipterix simplicella* (Stephens, 1834): E, ovipositor and 8th sternite; F, bursa copulatrix; G, *Ardania onobrychiella* (Zeller, 1849), male genitalia; H, *Casignetella heihensis* (Li et Zhang, 2000), male genitalia, <scale bars: 0.1 mm(A–D, G–H); 0.5 mm(E); 0.2 mm(F)>.

according to the territorial division adopted from the Catalogue of the Lepidoptera of Russia (Sinev 2008a).

In the taxonomic list, the novel records are marked with asterisks: for Krasnoyarskii Krai and/or the Republic of Khakassia by one symbol [*], novel records for Siberia with two symbols [**], species for the first time recorded on the territory of the Russia with three symbols [***], quarantine species, introduced with fruits and treated now as adventive species, with plus sign [+]. After abbreviation is used for the pheromone traps with the attractant for a particular insect species: pheromone trap with the attractant for *Grapholita molesta* (PHTGM), for *Dendrolimus sibiricus* (PHTDS), for *Frankliniella occidentalis* (PHTFO), for *Lymantria dispar* (PHTLD), for *Carposina "niponensis"* (PHTCN), and for *Ceratitis capitata* (PHTCC). Collectors: Evgeny N. Akulov (ENA), Natalia I. Kirichenko (NIK), Mikhail A. Ivanov (MAI).

The specimens are stored in the collection of the Krasnoyarsk branch of the All-Russian Center for Plant Quarantine, and the vouchers of *Coleophora gryphipennella* and *C. curictae* are stored in the collection Sukachev Institute of Forest SB RAS (Krasnoyarsk, Russia).

Systematic accounts

Superfamily Nepticuloidea
Family Nepticulidae

Bohemannia pulverosella (Stainton, 1849) **
(Figure 3A)

Material examined. 9 ♀, Krasnoyarskii Krai, Krasnoyarsk, V.M. Krutovsky Botanical Garden, orchard, PHTGM, 17, 24 v 2017 (ENA).

Bionomics. Leaf-miner of the *Malus* and *Pyrus* (Rosaceae) (Emmet 1979; Heath 1983). Only females were collected in south Siberia which suggests that the species is probably parthenogenetic in that area same as most European populations (Nieukerken et al 2016).

Remarks. Transholarctic species with disjunction in geographic range on the territory of Central Eurasia, it was introduced in North America (Nieukerken et al 2016). Russia: North-Western European part, Middle Volga Region and South Ural, and Primorskii Krai (as *piotra* Puplesis 1984) (Sinev 2008b). First record for Siberia (present article).

Superfamily Incurvarioidea
Family Adelidae

Nematopogon robertella (Clerck, 1759) *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", forest, at daytime, 15 vi 2017 (ENA).

Bionomics. Phytophagous species. In Europe, the larvae of younger age are leaf miner, and later they live in foliar case and feed on withered leaves on the litter in coniferous-deciduous forest. The foliar case maden from *Fagus sylvatica* (Fagaceae) was recorded in Switzerland (Fagaceae) (Lepiforum 2019) and *Vaccinium myrtillus* (Ericaceae) as a host plant in Europe (Kozlov 2016).

Remarks. Eurasian temperate species. Russia: Kola Peninsula, Karelia Republic, North-Western, North-Eastern and Central European parts, Middle Volga Region, Altaiskii Krai, Tuva Republic, and South Kuril Isl. (Kozlov 2008, 2016). First record for Krasnoyarskii Krai (present article).

Superfamily Tineoidea
Family Tineidae

Archinemapogon yildizae (Koçak, 1981) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, HTLD, viii 2017 (ENA).

Bionomics. Phytophagous species. Larvae feed in rotten wood affected by fungal hyphae and in bracket fungi (Polyporaceae) growing on deciduous trees.

Remarks. Eurasian temperate species. Russia: Kola Peninsula, Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, Middle and South Ural, Altaiskii Krai and Primorskii Krai (Baryshnikova 2008; Ponomarenko 2016). First record for Khakassia Republic (present article).

Nemapogon variatella (Clemens, 1859) *

Material examined. 1 ♀, Krasnoyarskii Krai, Krasnoyarsk, V.M. Krutovsky Botanical garden, orchard, PHTGM, 07 vi 2017 (ENA).

Bionomics. Phytophagous species. The larvae feed in decayed wood affected by fungal hyphae and in bracket fungi (Polyporaceae), and they also can develop in storages and in warehouses with grain and products of its processing (Emmet 1988).

Remarks. Transholarctic species. Russia: Kaliningradskaya Oblast, North-Western and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, Middle and South Ural, Altaiskii Krai, Irkutskaya Oblast, Zabaikalskii Krai, and Primorskii Krai (Baryshnikova 2008; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Monopis obviella ([Denis et Schiffermüller], 1775) **
(Figure 3B)

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, HTLD, viii 2017 (ENA).

Bionomics. Saprophagous species. The larvae feed on substrate of animal and plant (especially wood) origin (Emmet and Heath 1991).

Remarks. Euro-Siberian species. Russia: Kaliningradskaya Oblast, Central European part, Middle Volga and Volga-Don Regions, North Caucasus, and South Ural (Baryshnikova 2008). First record for Siberia (present article), Khakassia Republic is most eastern locality in distribution.

Monopis weaverella (Scott, 1858) *

Material examined. 1 ♀, 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 24 vi 2018 (ENA).

Bionomics. Saprophagous species. The larvae feed on substrate of animal origin (Heath and Emmet 1985).

Remarks. Transholarctic species with disjunction in geographic range in East Eurasia. Russia: Kola Peninsula, North-Western and Central European parts, Middle Volga, South Ural, North Enisej Region, Altaiskii Krai, and Primorskii Krai (Baryshnikova 2008; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Family Psychidae genus *Proutia*

Proutia betulina (Zeller, 1839) *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, Vet-luzhanka, private gardens' community «KZK-2», PHTGM, 26 vi 2018 (ENA).

Bionomics. Phytophagous species. In Europe, the larvae feed on lichens and decaying leaves living in case from pieces of plant such as leaves, bark, and so forth, live in deciduous forests on the bark of *Betula* spp. (Betulaceae) (Sinev 2016).

Remarks. Eurasian temperate species. Russia: Karelia Republic, North-Western and Central Regions, Middle Volga and Volga-Don Regions, North Caucasus, south of West Siberia, Tuva Republic, Buryatia Republic, Amurskaya Oblast, and Primorskii Krai (Sinev and Lovtsova 2008; Sinev 2016). First record for Krasnoyarskii Krai (present article).

Superfamily Gracillarioidea
Family Bucculatricidae

Bucculatrix artemisiella (Herrich-Schäffer, 1855) *

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Kubekovo settl., at daytime on herbaceous vegetation, 13 v 2017 (ENA).

Bionomics. Phytophagous species mining leaves on *Artemisia campestris* (Asteraceae) (Lepiforum 2019).

Remarks. Euro-Siberian species. Russia: Central European part, Middle Volga, Volga-Don and Low Volga Regions, Zabaikalskii Krai (Baryshnikova 2008). First record for Krasnoyarskii Krai (present article).

Bucculatrix bechsteinella (Beckstein et Scharfenberg, 1805) *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, market with import fruit production, PHTCN, 03 viii 2017 (ENA).

Bionomics. Phytophagous species, facultative leaf-miner and leaf-skeletonizer on *Crataegus*, *Malus*, *Pyrus*, and *Sorbus* (Rosaceae) (Emmet and Heath 1991).

Remarks. Euro-Siberian species. Russia: Kaliningradskaya Oblast, North-Western and Central European parts, Middle Volga, South Ural, and Buryatia Republic (Baryshnikova 2008). First record for Krasnoyarskii Krai (present article).

Bucculatrix nigricomella (Zeller, 1839) **
(Figure 3C)

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTLD, viii 2017 (ENA).

Bionomics. Phytophagous species, mining and skeletonizing the leaves on *Leucanthemum vulgare* and *Chrysanthemum leucanthemum* (Asteraceae) (Emmet and Heath 1991).

Remarks. Euro-Siberian species. Russia: Kaliningradskaya Oblast, Karelia Republic, Central European part, and Middle Volga (Baryshnikova 2008). First record for Siberia (present article), Khakassia Republic is the most eastern locality of the species distribution.

Bucculatrix pannonica Deschka, 1982 ***
(Figure 3D)

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", steppe slope, at daytime, 22 vi 2018 (ENA).

Bionomics. Phytophagous species, mining leaves on *Artemisia maritima* (Asteraceae) (Deschka 1982).

Remarks. Euro-Siberian species, in Europe known only from Austria, Hungary, Slovakia, and Ukraine (Deschka 1982; Baraniak 1996; Bidzilya and Budashkin 2017). First record for Russia and Siberia (present article). Krasnoyarskii Krai is the most eastern locality in the species range.

Superfamily Yponomeutoidea
Family Yponomeutidae

Swammerdamia glaucella Junnilainen, 2001 *

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 24 vi 2018 (ENA).

Remarks. East-Palearctic species, with geographic range included South Ural, Tuva Republic and Buryatia Republic in Russia, also South-East Kazakhstan, Mongolia (Junnilainen 2001), and South Korea (Sohn 2009). First record for Krasnoyarskii Krai (present article).

Family Argyrethiidae

Argyresthia brockeella (Hübner, 1813) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA); 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species. In Europe, the larvae feed in catkins and buds on birch (*Betula*) и alder (*Alnus*) (Betulaceae) (Agassiz 1987; Emmet 1996).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, Karelia Republic, North-Western, North-Eastern, and Central European parts, Middle Volga Region; Middle and South Ural, south of West Siberia, Irkutskaya Oblast, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), Sakhalin Isl., and Primorskii Krai (Sinev 2008b; Ponomarenko 2016). First record for Khakassia Republic and Krasnoyarskii Krai (present article).

Family Ypsolophidae

Ypsolopha asperella (Linnaeus, 1761) *

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 17 iv 2017 (MAI).

Bionomics. Phytophagous species, the larvae live in spinning a slight web and feed on the leaves of *Malus*, *Prunus*, *Crataegus*, and *Pyrus* (Rosaceae) (Emmet 1996).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, South Ural, Altai Republic, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), Sakhalin Isl., and Primorskii Krai (Sinev 2008b; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Ypsolopha dentella (Fabricius, 1775) *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, at light, 22 vii 2018 (MAI).

Bionomics. Phytophagous species, the larvae feed on *Lonicera* sp. and *Symphoricarpos rivularis* (Caprifoliaceae) (Agassiz 1983).

Remarks. Transholarctic species. Russia: Kaliningradskaya Oblast, North-Western, North-Eastern and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, South Ural, Altai Republic, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Sinev 2008b; Ponomarenko 2016; Huemer et al 2017). First record for Krasnoyarskii Krai (present article).

Ypsolopha falcella ([Denis et Schiffermüller], 1775) *

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 14 iv 2018 (ENA).

Bionomics. Phytophagous species, the larvae feed on *Lonicera xylosteum*, *L. carprifolium* (Caprifoliaceae) (Gershenson and Kozhevnikova 2013).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, North-Western and Central European parts, Middle Volga Region, Middle Ural, Altai Republic, Irkutskaya Oblast, Zabaikalskii Krai, Khabarovskii Krai (south), and Primorskii Krai (Sinev 2008b; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Family Glyphipterigidae

Glyphipterix magnatella Erschoff, 1877 *

Material examined. 2 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", birch & pine forest, at daytime, 22 vi 2018 (ENA).

Remarks. East-Palearctic. Russia: Altai Republic, Irkutskaya Oblast, and Primorskii Krai (Sinev 2008b; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Glyphipterix simpliciella (Stephens, 1834) **
(Figure 3E–F)

Material examined. 1 ♀, Krasnoyarskii Krai, Krasnoyarsk, V.M. Krutovsky Botanical Garden, orchard, PHTGM, 07 vi 2017 (ENA).

Bionomics. Phytophagous, carpophagous species, the larvae feed on the seeds of *Dactylis glomerata* and *Festuca* spp. (Poaceae).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, Altaiskii Krai (S.Yu. Sinev, pers. commun.), and Primorskii Krai (Sinev 2008b; Ponomarenko 2016). First record for Siberia (present article).

Superfamily Gelechioidea

Family Depressariidae

Agonopterix alstromeriana (Clerck, 1759) *

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl. vicinity, forest steppe, at light, 17 iv 2017 (MAI); 1 ♂, Krasnoyarsk, Vertluzhanka, private gardens' community «KZK-2», PHTGM, 24 v 2017 (ENA); 1 ♂, Berezovskii District, Berezovskii settl., at light, 14 iv 2018 (ENA).

Bionomics. Phytophagous, the larvae feed on hemlock (*Conium maculatum*, Apiaceae), rolling leaves, sometimes together with flowers and developing seeds (Berenbaum and Passoa 1983).

Remarks. Transholarctic species introduced in New Zealand (Winston et al 2014; Hayes 2019). Russia: Kaliningradskaya Oblast, Karelia Republic, North-Western, North-Eastern and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, Middle and South Ural, and Altai Republic (Lvovsky 2006, 2008). First record for Krasnoyarskii Krai (present article); this locality is the most eastern in territory of species existence in Russia.

***Agonopterix bipunctosa* (Curtis, 1850) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 20 viii 2015 (MAI); 15 ♂, same locality, 18 viii 2018 (EHA).

Bionomics. Phytophagous species, the larvae leaf-roller, feeding on *Centaurea scabiosa* and *Serratula tinctoria* (Asteraceae) (Lvovsky 2006).

Remarks. Euro-Siberian species. Russia: South Ural, south of West Siberia (Lvovsky 2006). First record for Krasnoyarskii Krai (present article), this locality is most eastern in the species range.

***Agonopterix multiplicella* (Erschoff, 1877) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, V.M. Krutovsky Botanical Garden, orchard, PHTGM, 07 vi 2017 (ENA); 1 ♂, Minusinskii district, Opytnoe pole settl., orchard, PHTGM, 15 v 2018 (ENA).

Bionomics. Phytophagous species, the larvae live and feed between webbed leaves of *Artemisia vulgaris* (Asteraceae) (Lvovsky 2006).

Remarks. Eurasian temperate species. Russia: North-Western and Central European parts, Middle Volga and Volga-Don Regions, South Ural, south of West Siberia, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Lvovsky 2008; 2016). First record for Krasnoyarskii Krai (present article).

***Agonopterix propinquella* (Treitschke, 1835) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species, facultative leaf-miner. The larvae of young ages mining the leaves of different Asteraceae (*Cirsium*, *Serratula*, *Carduus*, *Centaurea*, *Arctium*, *Cirsium*), the larvae of older ages feed in webbed leaves or in web under the leaf midrib (Lvovsky 2006).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, North-Western, North-Eastern and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, South Ural, south of West Siberia, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Lvovsky 2008). First record for Krasnoyarskii Krai (present article).

***Depressaria depressana* (Fabricius, 1775) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, at light, 05 x 2017 (MAI).

Bionomics. Anthophagous, rarely phytophagous species. The larvae live and feed in inflorescences and leaves of different Apiaceae (*Daucus*, *Pastinaca*, *Peucedanum*, *Pimpinella*, *Seseli*, and others) (Lvovsky 2006).

Remarks. Transpalearctic species. Russia: Kaliningradskaya Oblast, Karelia Republic, North-Western, North-Eastern and Central European parts, Middle Volga, Volga-Don and Low Volga Regions, North Caucasus, Middle and South Ural, south of West Siberia, Altai Republic, Zabaikalskii Krai, and Primorskii Krai (Lvovsky 2008, 2016). First record for Krasnoyarskii Krai (present article).

***Depressaria golovushkini* Lvovsky, 1995 ***

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, V.M. Krutovsky Botanical Garden, orchard, PHTCN, vii 2017 (ENA).

Bionomics. Phytophagous species. The larvae on *Diospyros* (Ebenaceae) in China (Lvovsky 2016).

Remarks. East-Palearctic species. Russia: Buryatia Republic, Zabaikalskii Krai, and Amurskaya Oblast (Lvovsky 2008, 2016). First record for Krasnoyarskii Krai (present article), this locality is most western in the species range.

Family Elachistidae

***Elachista adscitella* Stainton, 1851 ***

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTLD, vii 2017 (ENA).

Bionomics. Phytophagous species, the larvae are leaf-miners on *Deschampsia caespitosa*, *Brachypodium sylvaticum*, *Sesleria caerulea*, *Poa*, *Festuca altissima*, *F. gigantea*, *Milium effusum* (Poaceae) (Traugott-Olsen and Nielsen 1977).

Remarks. Eurasian temperate species, with disjunction in area in West Siberia. Russia: North-Western and Central European parts, North Caucasus, South Ural, Irkutskaya Oblast, and Primorskii Krai (Sinev 2008b, 2016). First record for Khakassia Republic (present article).

Family Scythrididae

***Scythris pudorinella* (Möschler, 1866) ***

Material examined. 1 ♀, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTDS, vii 2017 (ENA).

Remarks. Euro-Siberian species. Russia: Middle Volga, Volga-Don and Low Volga Regions, South Ural (Sinev 2008b, Nupponen 2009). First record for Khakassia Republic (present article), this locality is most eastern in the species range.

Family Chimabachidae

***Dasystoma salicella* (Hübner, 1796) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl. vicinity, forest steppe, at light, 17 iv 2017 (MAI).

Bionomics. Phytophagous species, the larvae are polyphagous and feed on different plants from Betulaceae (*Betula*), Salicaceae (*Salix caprea*), Ericaceae (*Rhododendron*, *Vaccinium myrtillus*), Rosaceae (*Rubus fruticosus*, *Filipendula ulmaria*, *Prunus spinosa*, *Potentilla anserina*), Rhamnaceae (*Rhamnus cathartica*), and Myricaceae (*Myrica gale*) (Emmet and Langmaid 2002; Lvovsky 2006).

Remarks. Transholarctic species. Russia: Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, Irkutskaya Oblast, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Lvovsky 2008, 2016). First record for Krasnoyarskii Krai (present article).

Family Oecophoridae

***Bisigna procerella* ([Denis et Schiffermüller], 1775) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", birch-pine forest, at daytime, 22 vi 2018 (ENA).

Bionomics. Phytophagous species, the larvae feed on lichens on the trunks of the trees, they were found on *Betula* (Betulaceae) and *Fagus* (Fagaceae) (Lvovsky 2006).

Remarks. Eurasian temperate species. Russia: North-Western and Central European parts, Middle Volga and Volga-Don Regions, South Ural, Altai Republic, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Lvovsky 2008, 2016). First record for Krasnoyarskii Krai (present article).

***Epicallima formosella* ([Denis et Schiffermüller], 1775) ***

Material examined. 1 ♀, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species, the larvae live and feed under dead bark, in Europe they were found on the *Malus domestica* (Rosaceae), *Robinia pseudacacia* (Fabaceae), and *Populus* (Salicaceae) (Emmet and Langmaid 2002; Lvovsky 2006).

Remarks. Transholarctic species, with disjunction in range in Eastern Eurasia. Russia: North-Western and Central European parts, Middle Volga and Volga-Don Regions, South Ural, and south of West

Siberia (Lvovsky 2008). First record for East Siberia (present article), Krasnoyarskii Krai is the most eastern locality in the species range.

***Endrosis sarcitrella* (Linnaeus, 1758) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA).

Bionomics. Saprophagous species, the larvae feed on remains of plant and animal origin and stored goods.

Remarks. Subcosmopolitan synanthropic species, introduced into different parts of the world. Russia: Kaliningradskaya Oblast, Kola Peninsula, Karelia Republic, North-Western, North-Eastern and Central European parts, Middle Volga and Volga-Don Regions, Middle and South Ural, Altaiskii Krai, Irkutskaya Oblast, Zabaikalskii Krai, South Yakutia, Kamchatka Peninsula, Khabarovskii Krai (south), and Primorskii Krai (Lvovsky 2008, 2016). First record for Krasnoyarskii Krai (present article).

Family Coleophoridae

***Coleophora gryphipennella* (Hübner, 1796) ***
(Figure 4A–D)

Material examined. 2 feeding larvae in the cases (Figure 4A–D), Krasnoyarskii Krai, Pogorelskiy bor, Arboretum SIF SB RAS, *Rosa* sp., 16 ix 2013 (NIK); one DNA barcoded larva: sample ID (NK253), process ID (ISSIK117-14), BIN (BOLD:AAE8826).

Bionomics. Oligophagous species on Rosaceae. In Europe, larvae develop on *Rosa acicularis*, *R. arkansana*, *R. canina*, *R. glauca*, *R. pendulina*, *R. rubiginosa*, *R. serafinii*, *R. soulieana*, *R. spinosissima*, *R. tomentosa* (Ellis 2019), occasionally on *Rubus caesius* and *R. corylifolius* (Hering 1957; Maček 1999). In the Netherlands, it was occasionally recorded on *Fragaria vesca* (van Roosmalen and Doorenweerd, 2015), and in Japan, on *R. rugosa* and *R. suavis* (Anikin 2016). In Siberia, feeding larvae were found exceptionally on *Rosa* sp.

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, North-Western European part, Middle Volga and Volga-Don regions, Irkutskaya Oblast, Kamchatka Peninsula, and Primorskii Krai (Anikin 2016). First record for Krasnoyarskii Krai (present article).

The DNA barcode of the specimen from Siberia matches that of *C. gryphipennella* specimens from Central and Northern Europe

(Netherlands, Belgium, France, Italy, Austria, Germany, Bulgaria, Finland, Norway, Denmark), with 97.70–98.88% min-max similarity. The nearest neighbor of the Siberian specimen, with 98.88% similarity, is the specimen from Austria (process ID: PHLAI641-13.COI-5P, GenBank Accession: KM573253) (Huemer et al 2014). All DNA barcodes of *C. gryphipennella* have unique BIN (BOLD:AAE8826), and the Siberian specimen correspond to this BIN. Keeping in mind high genetic proximity of the Siberian specimen and European specimens identified as *C. gryphipennella* and their possession to one BIN, the larva feeding on *Rosa* sp. in Krasnoyarskii Krai is assigned to *C. gryphipennella*. Additional sampling and rearing of adults in the region would be needed to morphologically confirm the species.

Coleophora curictae* Baldizzone, 2016 **
(Figure 5A–H)

Material examined. Two leaves with mines (Figure 5A–B) and 3 feeding larvae in the cases (Figure 5C–H), Krasnoyarskii Krai, Krasnoyarsk, Red Square, trees along the road, *Ulmus pumila*, 12 vi 2013 (NIK); one DNA barcoded larva: sample ID (NK241), process ID (ISSIK105-14), BIN (BOLD:AAM2030).

Bionomics. Neither biology nor host plants are known for the species in Europe (Baldizzone 2016). However, Baldizzone (2016) suggested that it may probably feed on *Quercus*. In Siberia, we found feeding larva on *Ulmus pumila* L. (Ulmaceae) (Figure 5A–D). Based on our observations in Krasnoyarsk, larval development takes place in June (starting in late May and probably lasting till sometime in July). Larva feeds moving on the leaf making oval blotch mines; several mines per leaf made by one larva can be recorded (Figure 5A–D). When the leaf is significantly mined, the larva moves to a neighboring leaf and continues feeding. Larva has rough pistol case with swirling rear part, about 8 mm in length, with a mouth angle of 15°–25° standing obliquely on the leaf. The case of a younger larva is brownish-silvery, later with a more extended mouth part (Figure 5E–F).

Remarks. Euro-Siberian species. First record for Russia based on our sampling in Krasnoyarskii Krai (present article).

The specimen from Siberia was identified through DNA barcoding. It showed 98.58% similarity with an adult moth sampled in




Figure 4. Biology of *Coleophora gryphipennella* (Hübner, 1796) in Krasnoyarskii Krai.: A, larva feeding on *Rosa* leaf; B, the leaf mine; C–D, the larval case. <scale bars: 12 mm (A); 5 mm (B); 6 mm (C–D)>.

Tyrol (Austria), process ID: LEATI273-15.COI-5P (Huemmer and Hebert 2016). Based on a high genetic similarity between the Siberian and Austrian specimens, that the species feeding in Siberia on *Ulmus pumila* is *Coleophora curictae*. Additional sampling and rearing of adults in the region would be needed to morphologically confirm the species.

Ardania onobrychiella (Zeller, 1849) **
(Figure 3G)

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species on *Astragalus arenarius*, *Onobrychis vicifolia*, *Hyppocrepis comosa* (Fabaceae) (Falkovitsh 2006).

Remarks. Euro-Siberian species with vast disjunction in range. Russia: Kaliningradskaya Oblast (Anikin 2008). First record for Siberia with only locality in Krasnoyarskii Krai (present article).

Ardania sergiella Falkovitsh 1979 *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoturanskii District, Novaya Syda settl., steppes slope, 12 vi 2018 (ENA).

Bionomics. Phytophagous species on *Potentilla cinerea* (Rosaceae).

Remarks. South-Siberian species. Russia: North Caucasus, Altai Republic, Tuva Republic, Zabaikalskii Krai (Anikin 2008). First record for Krasnoyarskii Krai (present article).

Damophila alcyonipennella (Kollar, 1832) *

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl. vicinity, forest steppe, at light, 17 iv 2017 (MAI); 1 ♂, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA); 1 ♂, Emeljanovskii District, Krasnoyarsk Reservoir, Bay of Biryusa 30 vi 2018 (ENA).

Bionomics. Phytophagous species, feeds on inflorescences on *Trifolium* (Fabaceae) (Falkovitsh 2006).

Remarks. Transpalearctic species introduced into New Zealand and Australia. Russia: Kaliningradskaya Oblast, North-Western and Central European parts, Middle Volga, Low Volga Regions and Volgo-Don Regions, North Caucasus, South Ural, south of West Siberia, Tuva Republic, Irkutskaya Oblast, Yakutia Republic (north-east), Khabarovskii Krai (south), Sakhalin Isl., and Primorskii




Figure 5. Biology of *Coleophora curictae* Baldizzzone, 2016 in Krasnoyarskii Krai.: A, mines on *Ulmus pumila* L.; C–D, larva feeding in the mine not entirely leaving the case attached to the mine; E, the case of a young larva (lateral view); F, the case of the older larva (lateral view); G–H, larva (G, dorsal and H, lateral views). <scale bars: 20 mm (A); 8 mm (B); 7 mm (C–D); 3 mm (E); 4 mm (F); 2 mm (G–H)>.

Krai (Anikin 2016). First record for Krasnoyarskii Krai (present article).

***Damophila trifolii* Curtis, 1832 ***

Material examined. 1 ♀, 1 ♂, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA); 1 ♀, Krasnoyarskii Krai, Emeljanovskii District, Minino settl. vicinity, forest steppe, at light, 17 iv 2017 (MAI); 1 ♀, Emeljanovskii District, Krasnoyarsk Reservoir, Biryusa Bay 30 vi 2018 (ENA); 1 ♀, Krasnoyarsk, V.M. Krutovsky Botanical Garden, orchard, at daytime on Fabaceae, 11 vii 2018 (ENA); 1 ♂, Krasnoyarsk, at light, 04 viii 2018 (ENA).

Bionomics. Carpophagous species feeds on seeds of *Melilotus officinalis*, *M. altissimus* (Falkovitsh 2006), and probably on *Trifolium* (Fabaceae).

Remarks. Transholarctic species. Russia: Central European part, Middle Volga, Volga-Don Regions, North Caucasus, Ural, south of West Siberia, Altai Republic, Irkutskaya Oblast, Zabaikalskii Krai, and Primorskii Krai (Anikin 2016). First record for Khakassia Republic and Krasnoyarskii Krai (present article).

***Multicoloria vibicigerella* (Zeller, 1839) ***

Material examined. 1 ♀, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species, the larvae mine leaves *Achillea millefolium*, *A. ptarmica*, and *Artemisia campestris* (Asteraceae) (Anikin 2004).

Remarks. Transpalearctic species. Russia: North-west and Central European parts, Middle Volga, Low Volga, Volga-Don Regions, North Caucasus, South Ural, Altai Republic, Tuva Republic, questionably Zabaikalskii Krai, and Primorskii Krai (Anikin 2008, 2016). First record for Khakassia Republic (present article).

***Ecebalia hsiaolingensis* (Toll, 1942) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, at light, 11 ix 2017 (ENA); 3 ♂, Berezovskii District, Berezovskii settl., at light, 12, 28 viii 2018 (ENA); 3 ♂, Emeljanovskii District, Minino settl., steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species, the larvae feed on inflorescence of *Aster* (Asteraceae) (Anikin 2016).

Remarks. East-Palearctic species. Russia: south of West Siberia, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai, Magadanskaya Oblast, and Primorskii Krai (Anikin 2016). First record for Krasnoyarskii Krai (present article).

***Ecebalia therinella* (Tengström, 1848) ***

Material examined. 2 ♀, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous, carpophagous species, the larvae feed on seeds of *Fallopia convolvulus* (Polygonaceae) (Falkovitsh 2006) or mine the leaves of *Carlina*, *Carduus*, and *Cirsium* (Asteraceae) (Anikin 2016).

Remarks. Eurasian temperate species. Russia: Kaliningradskaya Oblast, North-Western and Central European parts, Middle Volga, Low Volga, Volga-Don Regions, North Caucasus, South Ural, south of West Siberia, Irkutskaya Oblast, Zabaikalskii Krai, Yuakutia Republic, Amurskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Anikin 2016). First record for Khakassia Republic (present article).

***Ecebalia versurella* (Zeller, 1849) ***

Material examined. 2 ♀, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species on *Chenopodium*, *Atriplex*, *Halimione*, and *Amaranthus* (Amaranthaceae) (Falkovitsh 2006).

Remarks. Subcosmopolitan species unknown in Africa and Australian Region. Russia: Kaliningradskaya Oblast, Karelia Republic, Central European part, Middle Volga, Low Volga, Volga-Don Regions, North Caucasus, South Ural, Altai Republic, Khabarovskii Krai (south), and Primorskii Krai (north) (Anikin 2008, 2016). First record for Khakassia Republic (present article).

***Casignetella burmanni* (Toll, 1952) ***

Material examined. 2 ♀, Khakassia Republic, Shirinskii District, Chjerno Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species, the larvae mine leaves of *Gypsophila fastigiata* and *G. repens* (Caryophyllaceae) (Falkovitsh 2006).

Remarks. Euro-Siberian species. Russia: Altaiskii Krai (Anikin 2008). First record for Khakassia Republic (present article).

***Casignetella heihensis* (Li et Zhang, 2000) ** (Figure 3H)**

Material examined. 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., steppe, at light, 18 viii 2018 (ENA).

Bionomics. Phytophagous species on *Artemisia* (Asteraceae).

Remarks. East-Palearctic species. Russia: Amurskaya Oblast and Primorskii Krai (Anikin 2016). First record for Siberia (present article). Krasnoyarskii Krai is the most western locality in the species distribution.

***Casignetella striatipennella* (Nylander in Tengström, [1848]) ***

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA).

Bionomics. Carpophagous species feed on seeds of *Stellaria*, *Cerastium*, *Moehringia*, and *Myosoton* (Caryophyllaceae) (Falkovitsh 2006).

Remarks. Eurasian temperate species. Russia: Kola Peninsula, Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, south of West Siberia, Altaiskii Krai, Altai Republic, Magadanskaya Oblast, Khabarovskii Krai (south), and Primorskii Krai (Anikin 2008, 2016). First record for Krasnoyarskii Krai (present article).

Family Momphidae

***Mompha langiella* (Hübner, 1796) ***

Material examined. 1 ♀, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at daytime, 28 v 2017 (ENA).

Bionomics. Phytophagous species, the larvae are leaf-miners on *Circaea lutetiana*, *C. alpina*, *Epilobium hirsutum*, *E. montanum*, *E. parviflorum*, *Chamaenorium angustifolium*, and *Oenothera biennis* (Onagraceae) (Koster and Sinev 2003).

Remarks. Euro-Siberian species. Russia: Kaliningradskaya Oblast, Central European part, Middle Volga Region (Sinev 2008b), and Altaiskii Krai (Irisova et al 2014). First record for Krasnoyarskii Krai (present article).

Family Cosmopterigidae

***Pancalia hexachrysa* (Meyrick, 1935) ** (Figure 6A)**

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", steppe slope, at daytime, 22 vi 2018 (ENA).

Remarks. East-Palearctic species. Russia: Primorskii Krai (Sinev 2008b, 2016). First record for Siberia (present article). Krasnoyarskii Krai is the most western locality in the species distribution.

***Cosmopterix sibirica* Sinev, 1985 ***

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA).

Remarks. Euro-Siberian species. Russia: Middle Volga Region, South Ural, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai (Sinev 2008b). First record for Krasnoyarskii Krai (present article).

Eteobalea serratella (Treitschke, 1833) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species, the larvae are stem-miners and root-miners on *Linaria vulgaris*, *L. genistifolia* ssp. *euxina* (Plantaginaceae) (Koster and Sinev 2003).

Remarks. Euro-Siberian species. Russia: Middle Volga, Volga-Don and Low Volga Regions, North Caucasus, South Ural, and Zabaikalskii Krai (Sinev 2008b). First record for Khakassia Republic (present article).

Family Gelechiidae

Metzneria metzneriella (Stainton, 1851) *

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA).

Bionomics. Carpophagous species. The larvae feed on seeds of *Serratula tinctoria*, *Centaurea nigra*, *C. jacea*, *C. scabiosa* (Asteraceae) (Elsner et al 1999; Bland, Emmet et al 2002).

Remarks. Eurasian temperate species. Russia: Karelia Republic, North-Western and Central European parts, Middle Volga, Volga-Don and Low Volga Regions, North Caucasus, South Ural, Zabaikalskii Krai, Amurskaya Oblast, and Khabarovskii Krai (south) (Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Argolamprotes micella ([Denis et Schiffermüller], 1775) *

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk: Studgorodok, private gardens' community «Pobeda», orchard, PHTGM, 11 vii 2018 (ENA).

Bionomics. Phytophagous species. The larvae feed on flower buds, shoots by boring them, and on leaves skeletonizing the latter on *Rubus idaeus*, *R. fruticosus*, and *R. caesius* (Rosaceae) (Elsner et al 1999; Bland, Emmet et al 2002).

Remarks. Eurasian temperate species. Russia: North-West and Central European parts, Middle Volga Region, Middle and South Ural, Irkutskaya Oblast, Zabaikalskii Krai, Amurskaya Oblast,




Figure 6. Male genitalia: A, *Pancalia hexachrysa* (Meyrick, 1935), lateral view; B, *Tuta absoluta* (Meyrick, 1917), ventral view; C, *Caryocolum repentis* (Hübner et Luquet, 1992), ventral view; D-E, *Carposina sasakii* (Matsumura, 1900), D, ventral view; E, aedeagus. <scale bars: 0.1 mm>.

Khabarovskii Krai (south), Sakhalin Isl., Kuril Isl., and Primorskii Krai (Junnilainen et al 2010; Ponomarenko 2016). First record for Krasnoyarskii Krai (present article).

Monochroa simplicella (Lienig et Zeller, 1846) *

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 10 vi 2017 (ENA).

Bionomics. Phytophagous species. The larvae on *Juncus* sp. (Juncaceae) and *Eriophorum* sp. (Cyperaceae) (Elsner et al 1999).

Remarks. Euro-Siberian species. Russia: Central European part, South Ural, Altai Republic, and Irkutskaya oblast (Ponomarenko 2008; Bidzilya 2009; Junnilainen et al 2010). First record for Krasnoyarskii Krai (present article).

Chionodes tragicella (Heyden, 1865) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTDS, vii 2017 (ENA).

Bionomics. Phytophagous species. The larvae live under the scales of bark and feed on the needles of *Larix decidua* (Pinaceae) (Huemer and Karsholt 1999).

Remarks. Euro-Siberian species. Russia: Karelia Republic, Volga-Don Region, South Ural, Tuva and Buryatia Republics, and Zabaikalskii Krai (Anikin et al 1999, 2017; Bidzilya et al 1998; Ponomarenko 2008; Bidzilya 2009; Junnilainen et al 2010; Kutenkova et al 2015). First record for Khakassia Republic (present article).

Aroga controvalva Li et Zheng, 1998 *

Material examined. 4 ♂, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA); 1 ♂, Krasnoyarskii Krai, Emeljanovskii District, Minino settl., forest steppe, at light, 18 viii 2018 (ENA).

Remarks. East-Palearctic species. Russia: Zabaikalskii Krai (Kostjuk et al 1994, as *A. mesostrepta* (Meyrick 1932); Bidzilya et al 1998, as *A. trilineella* (Chambers 1877); Bidzilya 2009). First record for Khakassia Republic and Krasnoyarskii Krai (present article).

Filatima incomptella (Herrich-Schaffer, 1854) *

Material examined. 1 ♂, Krasnoyarskii Krai, Berezovskii District, Berezovskii settl., at light, 24 vi 2018 (ENA).

Bionomics. Phytophagous species. The larvae feed on leaves of *Salix caprea*, *S. aurita*, *S. bicolor*, *S. repens* (Salicaceae) (Huemer and Karsholt 1999).

Remarks. Euro-Siberian species. Russia: Karelia Republic, North-Western and Central European parts, Middle-Volga, Volga-Don and Low Volga Regions, South Ural, and Zabaikalskii Krai (Junnilainen et al 2010; Ponomarenko 2016; Ismagilov 2018). First record for Krasnoyarskii Krai (present article).

Gnorimoschema epithymella (Staudinger, 1859) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTLD, vii 2017 (ENA).

Bionomics. Phytophagous species. The larvae bore lower stems and mine the leaves of *Solidago virgaurea* и *Aster alpinus* (Asteraceae) (Huemer and Karsholt 1999).

Remarks. Euro-Siberian species. Russia: Kola Peninsula, Central European part, Middle-Volga, Volga-Don and Low Volga Regions, North Caucasus, and Zabaikalskii Krai (Ponomarenko 2008). First record for Khakassia Republic (present article).

Gnorimoschema jalavai Povolný, 1994 *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA).

Remarks. East-Palearctic species. Russia: Altai Republic, Tuva Republic, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai,

and Chukotian Region (Ponomarenko 2008). First record for Khakassia Republic (present article).

Tuta absoluta (Meyrick, 1917) ‡
(Figure 6B)

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, market with import fruit production, PHTCC, 24 vii 2019 (ENA).

Bionomics. Phytophagous species, pest of different Solanaceae, quarantine object for Russia. The larvae feed on different parts of host plant and have different feeding habits from leaf-mining to boring into buds, stalks, and fruits (Huemer and Karsholt 1999).

Remarks. Widely distributed species originated from South America and introduced into Europe, Africa, Nearest East, Asia Minor, and India. Invasive in Russia: Low Volga Region, North Caucasus, South Ural; for the first time recorded in Krasnoyarskii Krai (present article).

It is the object of the external quarantine in Russia, included into the List 1 "Quarantine objects absent in the territory of the Russian Federation" in the site of Federal Service for Veterinary and Phytosanitary Surveillance (Rosselkhoz nadzor 2019). Originated from South America, the tomato moth was recorded for the first time in Russia in North Caucasus (Krasnodarskii Krai) in 2010 (Izhevsky et al 2011). Later, the pest was found in other regions of North Caucasus (Adygea and Dagestan Republics) and in South Ural (Bashkortostan Republic) (Zhimerikin and Mironova 2012). Recently, it was captured in the Low Volga Region (Astrakhanskaya Oblast) (Medvedeva 2018). All these records are probably due to accidental movement of the pest with the imported vegetables, seeds, or seedlings. The data confirming naturalization of *T. absoluta* in these regions are absent. However, this potential pest could be able to settle down in the southern regions with low sum of the negative temperatures in winter, low soil freezing depth, and sufficient snow cover, as was shown in the experiments in south Ukraine (Klechkovsky et al 2014).

Agonochaetia Ivovskyi Bidzilya, 2001 *

Material examined. 3 ♂, Krasnoyarskii Krai, Krasnoyarsk, Nature reserve "Stolby", steppe slope, at daytime, 22 vi 2018 (ENA).

Remarks. South-Siberian species. Russia: Altai Republic (Bidzilya 2001). First record for Krasnoyarskii Krai (present article).

Caryocolum fischerella (Treitschke, 1833) **

Material examined. 2 ♀, Krasnoyarskii Krai, Krasnoyarsk, greenhouse complex, PHTFO, viii 2018 (ENA).

Bionomics. Phytophagous species. The larvae live and feed on spined terminal shoots and leaves on *Saponaria officinalis*, *S. oymoides* (Caryophyllaceae) (Huemer and Karsholt 1999).

Remarks. Euro-Siberian species. Russia: North-Western and Central European parts, Middle Volga, and Volga-Don Regions (Ponomarenko 2008). First record for Siberia (present article). Krasnoyarskii Krai is the most eastern locality in geographic range of this species.

Caryocolum repentis Huemer et Luquet, 1992 **
(Figure 6C)

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTLD, viii 2017 (ENA).

Bionomics. Phytophagous species. The larvae are facultative leaf-miner, mining the leaves in young ages, later they live and feed in pod-like dwelling, madden on the terminal part of shoots on *Gypsophila repens* (Caryophyllaceae) (Huemer and Karsholt 1999).

Remarks. Euro-Siberian species. Russia: South Ural (Junnilainen et al 2010). First record for Siberia (present article). Khakassia Republic is the most eastern locality in geographic range of this species.

Pseudotelphusa paripunctella (Thunberg, 1794) *

Material examined. 1 ♂, 1 ♀, Khakassia Republic, Shirinskii District, Tuimskoe forestry, PHTDS, viii 2017 (ENA).

Bionomics. Phytophagous species. The larvae feed on *Betula* spp. (Betulaceae), *Fagus* и *Quercus* (Fagaceae), *Myrica gale* (Myricaceae), *Hippophae rhamnoides* (Elaeagnaceae) (Sattler 1980).

Remarks. Eurasian temperate species. Russia: Kola Peninsula, Karelia Republic, North-Western and Central European parts, Middle Volga and Volga-Don Regions, North Caucasus, Middle and South Ural, south of West Siberia, Irkutskaya Oblast, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), Kuril Isl., and Primorskii Krai (Ponomarenko 2016). First record for Khakassia Republic (present article).

Anacamptis temerella (Lienig et Zeller, 1846) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species. The larvae live and feed on spinning terminal shoots and leaves of *Salix lapponum*, *S. caprea*, *S. repens*, and *S. phylicifolia* (Salicaceae) (Elsner et al 1999; Savela 2001; Bland et al 2002).

Remarks. Eurasian temperate species. Russia: North-Western and North-Eastern European parts, Middle Volga and Volgo-Don Regions, North Caucasus, Buryatia Republic, Zabaikalskii Krai, Amurskaya Oblast, Sakhalin Isl., and Primorskii Krai (Ponomarenko 2016). First record for Khakassia Republic (present article).

Dichomeris rasilella (Herrich-Schaffer, 1854) *

Material examined. 1 ♂, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species. The larvae feed on *Artemisia vulgaris*, *A. pontica*, *Acosta rhenana* и *Centaurea* spp. (Asteraceae) (Ponomarenko 1997; Elsner et al 1999; Savela 2001).

Remarks. Eurasian temperate species. Russia: Central European part, Volga-Don and Low Volga Regions, North Caucasus, South Ural, Altai Republic, Tuva Republic, Irkutskaya Oblast, Buryatia Republic, Zabaikalskii Krai, Amurskaya Oblast, Khabarovskii Krai (south), Kuril Isl., and Primorskii Krai (Ponomarenko 2008). First record for Khakassia Republic (present article).

Acanthophila latipennella (Rebel, 1937) *

Material examined. 1 ♀, Khakassia Republic, Shirinskii District, Chjernoje Ozero vil., at light, 16 vii 2018 (ENA).

Bionomics. Phytophagous species. The larvae feed on *Picea abies* (Pinaceae) (Elsner et al 1999).

Remarks. Eurasian temperate species. Russia: North-Western, North-Eastern and Central European parts, South Ural, Altaiskii Krai, Irkutskaya Oblast, Zabaikalskii Krai, Sakhalin Isl., and Primorskii Krai (Ponomarenko and Omelko 2003; Junnilainen et al 2010; Ponomarenko 2016; Ismagilov 2018). First record for Khakassia Republic (present article).

Superfamily Copromorphaidea
Family Carposinidae

Carposina sasakii Matsumura, 1900 ‡

(Figure 6D–E)

Carposina niponensis auct. (nec Walsingham 1900), error identification.

Material examined. 1 ♂, Krasnoyarskii Krai, Krasnoyarsk, market with import fruit production, PHTCN, 01 viii 2015 (ENA).

Bionomics. Phytophagous species. The larvae feed in fruits of different Rosaceae (*Prunus* spp., *Malus* spp., *Pyrus* spp.).

Remarks. The present species range includes East Asia, from where the species originated, and North America (Canada, east and south of the United States), where it was introduced and naturalized (North American Moth Photographers Group 2019). Russia: Amurskaya Oblast, Khabarovskii Krai (south), Kuril Isl., and Primorskii Krai. In Krasnoyarskii Krai, the species was intercepted in the pheromone trap in the market for the first time, suggesting introduction of the species with imported fruits.

The species, known as peach fruit moth, is the pest of Rosaceae in East Asia and the object of internal (in Russia) and external (in Europe) quarantine. Despite the series of scientific publications devoted to the morphology of *Carposina sasakii* (Diakonoff 1989; Ponomarenko 1999; Nasu et al 2010), the species has been constantly confused with *C. niponensis* Walsingham 1900 and both species names were erroneously treated as synonyms. Thus, the *Carposina niponensis* hitherto is mentioned instead of *C. sasakii* in handbooks on the pests (Savotikov and Smetnik 1995), and in list 2, "Quarantine objects restrictedly distributed in the territory of the Russian Federation" on official site of Federal Service for Veterinary and Phytosanitary Surveillance in Russia (Rosselkhoz nadzor: Normativnye dokumenty 2019). Also the "*Carposina niponensis* Walsingham" is erroneously indicated as the other scientific name for *C. sasakii* in the website of CABI (2019).

The lectotype of the species *C. sasakii* Matsumura 1900 was designated by Razowski and Kumata (1985). Detailed redescription of both species, *C. sasakii* and *C. niponensis*, with illustrations of the type material (adults and genitalia) was provided by Diakonoff (1989). In the latter, *C. sasakii* and *C. niponensis* are treated as two different species that are proven by their distinctive morphology. Therewith, *C. sasakii* is widely distributed in East Asia and well known mainly as a pest of *Prunus persica*, and other fruit trees of Rosaceae, whereas the only specimen of *C. niponensis* (Holotype) was known for a long time until the species was rediscovered by capturing two specimens in the pheromone trap for *C. sasakii* in Wakayama and Oita Prefectures in 2008 (Japan) (Nasu et al 2010).

Discussion

As a result of our samplings, in total 62 species from 42 genera and 18 Microlepidopteran families were added to the faunistic list of Krasnoyarskii Krai and Khakassia Republic. The two quarantine species intercepted in Krasnoyarsk are not counted because there are no data confirming their establishment in the region. Forty-four out of 62 species are new for Krasnoyarskii Krai, 22 species new for Khakassia Republic, 18 genera and 3 families (Glyptopterigidae, Chimabachidae and Oecophoridae) are registered for studied region for the first time.

Two species, i.e., *Bucculatrix pannonica* (Bucculatricidae) and *Coleophora curictae* (Coleophoridae), collected on the territory of the city Krasnoyarsk and the suburban area, are new records for Russia. Taking into account the latter, overall eleven species are novel for Siberia. Among them, there are the representatives of nine families: Nepticulidae (*Bohemannia pulverosella*), Tineidae (*Monopis obviella*), Bucculatricidae (*Bucculatrix nigricomella*), Glyptopterigidae (*Glyptopterix simpliciella*), Coleophoridae (*Ardania onobrychiella*, *Casignetella heihensis*), Cosmopterigidae (*Pancalia hexachrysa*), and Gelechiidae (*Caryocolum fischerella*, *C. repens*).

The two species detected for the first time in Siberia (the tomato leaf-miner, *Tuta absoluta*, and the peach fruit moth, *Carposina sasakii*) are known as serious agricultural pests. *T. absoluta*,

originally known from South America, is a quarantine pest in Russia, unintentionally introduced to the European part of the country, in particular to Krasnodarskii Krai. Native to East Asia, *C. sasakii* has restricted distribution in Russia (with a primary range in the Russian Far East), and thus, it has a quarantine status for the rest of country. Both species were intercepted on the fruit markets in Krasnoyarsk using pheromone traps. They could be introduced into the region accidentally, with fruit crops from the regions with those Krasnoyarsk Krai has trade relations (Central and East Asia, Europe). The records of these two species in South Siberia are nominative and should be treated with a caution. The chances for *T. absoluta* and *C. sasakii* to establish the populations in Siberia are low keeping in mind unfavorable climatic conditions for these species originating from warmer areas. Nevertheless, the question about naturalization of these species on the southern territories of Krasnoyarsk Krai would require further monitoring programs.

Conclusions

Our study highlights the importance of regional inventories for defining faunal composition, clarifying species ranges, and revealing unwelcome guests—the invasive and quarantine species. As shown in this article, gathering the diverse collection on Microlepidoptera in South Siberia comprising exclusively of novel faunistic and geographic records was only possible thanks to the involvement of various sampling techniques allowing to cover different taxonomic groups of the moths. Besides continuing exploring the fauna of Microlepidoptera in South Siberia, our future goals will be to specify trophic associations and geographic ranges of the indigenous species and to explore the potential of nonnative pestiferous species, that might be accidentally introduced to the region through various pathways, to establish and spread in the region.

Conflict of interest

The authors declare that there is no conflict of interest.

Acknowledgments

The authors thank M. Ivanov (Krasnoyarsk) for helping with sampling on the territory of Krasnoyarskii Krai, A. Knorre (Nature reserve «Stolby», Krasnoyarsk) for her cooperation, I. Mikhailova and Yu. Baranchikov (SIF SB RAS, Krasnoyarsk), respectively, for help with mapping and for support at different stages of the study. Special thanks to the taxonomists for confirming of species identification from some families: V. Anikin (Coleophoridae), A. Lvovsky (Depressariidae), Yu. Lovtsova (Psychidae), S. Sinev (Nepticulidae, Glyphipterigidae, Elachistidae, Momphidae, Cosmopterigidae), and S. Baryshnikova (Bucculatricidae).

This work was supported by grants of the Russian Foundation for Basic Research, Russia (projects numbers 15-29-02645-ofi_m and 18-04-00944), LE STUDIUM® fellowship program, Institute for advanced studies—Loire Valley (Orléans, France) and Cost Action FP1401—A global network of nurseries as early warning system against alien tree pests (Global Warning).

References

- Agassiz DJL. 1983. Microlepidoptera — a review of the year 1982. *The Entomologist's Record and Journal of Variation* 95:187–195.
- Agassiz DJL. 1987. The British Argylethiinae and Yponomeutinae. *Proceedings and Transactions of the British Entomological and Natural History Society* 20 (1):1–26.
- Akulov EN, Belyakova OV, Kirichenko NI. 2013. Detection of the oriental fruit moth *Grapholita molesta* in the south part of Siberia. *Plant Protection and Quarantine* 10:34–37 [in Russian].
- Akulov E, Kirichenko N, Petko V. 2014. Non-targeted species of *Grapholita* (Lepidoptera, Tortricidae) attracted to synthetic pheromone of Oriental fruit moth in southern Siberia. *Plant Health. Research and Practice* 2 (8):31–50 [in Russian].
- Akulov EN, Kirichenko NI. 2014. Mass catches of the Manchurian moth in the south of Siberia. *Plant Protection and Quarantine* 10:36–40 [in Russian].
- Akulov EN, Kirichenko NI, Takhtuev AS. 2015. A dangerous solanaceous pest, *Leucinodes orbonalis*, was found in imported products in Russia. *Plant Protection and Quarantine* 8:36–38 [in Russian].
- Akulov EN, Kirichenko NI, Ponomarenko MG. 2018. Contribution to the Microlepidoptera fauna of the South of Krasnoyarsk Territory and the Republic of Khakassia. *Entomological Review* 98 (1):49–75. <https://doi.org/10.1134/S0013873818010074>.
- Anikin VV. 2004. Food connections of cover moths (Lepidoptera, Coleophoridae) in legume (Fabaceae) and complex-colored (Asteraceae) families on the territory of the Lower Volga. *Bulletin of the Botanical garden of Saratov state University* 3 (1):61–66.
- Anikin VV. 2008. Family Coleophoridae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. pp. 69–82 [in Russian].
- Anikin VV. 2016. Fam. Coleophoridae. In: Beljaev EA, editor. *Annotated catalogue of the insects of Russian Far East*, vol. II. Lepidoptera. Vladivostok: Dalnauka. pp. 97–105 [in Russian].
- Anikin VV, Sachkov SA, Zolotuhin VV. 1999. "Fauna Lepidopterologica Volgo-uralensis" 150 years later: changes and additions. Part 3. Coleophoridae, Gelechiidae, Symmocidae and Holcopogonidae (Insecta, Lepidoptera). *Atalanta* 29 (1/4):295–336.
- Anikin VV, Sachkov SA, Zolotuhin VV. 2017. "Fauna Lepidopterologica Volgo-uralensis": from P. Pallas to present days. *Proceedings of the Museum Witt Munich. Munich—Vilnius* 7:1–696.
- Baldizzone G. 2016. *Coleophora curictae* Baldizzone: a new species of the *C. zelleriella* Heinemann, 1854 group. Contribution to the knowledge of the Coleophoridae. CXXXVI (Lepidoptera: Coleophoridae). *SHILAP Revista de Lepidopterología* 44 (175):455–462.
- Baraniak E. 1996. Bucculatricidae. In: Karsholt O, Razowski J, editors. *The Lepidoptera of Europe. A distributional checklist*. Stenstrup: Apollo Books. pp. 47–48.
- Baryshnikova SV. 2008. Family Tineidae. Family Bucculatricidae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St. Petersburg, Moscow: KMK Scientific Press Ltd. pp. 27–32. 37–38. [In Russian].
- Berenbaum M, Passoa S. 1983. Notes on the biology of *Agonopterix alstroemeriana* (Clerck), with descriptions of the immature stages. *Journal of the Lepidopterists' Society* 37 (1):38–45.
- Bidzilya AV. 2001. Data to the distribution of gelechiid moths (Lepidoptera: Gelechiidae) in Southern Siberia. Report I (2002) *Izvestiya Kharkovskogo entomologicheskogo Obshchestva* 9 (1-2):64–72 [In Russian].
- Bidzilya OV. 2009. On the distribution of gelechiid moths (Lepidoptera: Gelechiidae) in Siberia. Contribution 3. *Proceedings of Zoological Museum Kiev Taras Shevchenko National University* 5:3–13.
- Bidzilya OV, Budashkin Yul. 2017. New records of Lepidoptera from Ukraine and description of a new species of *Caloptilia* Hübner, 1825 (Lepidoptera, Gracillariidae) from the mountains of Crimea. *Nota Lepidopterologica* 40 (2):5–21 [In Russian].
- Bidzilya OV, Budashkin Yul, Kostjuk IYu. 1998. Additions to the Fauna of Microlepidoptera of Transbaikalia. *Journal of Ukrainian Entomological Society* 4:33–63 [In Russian].
- Bland KP, Emmet AM, Heckford RJ, et al. 2002. Anomologinae. In: Emmet AM, editor. Langmaid JR, editor. *The moths and Butterflies of Great Britain and Ireland*, vol. 4. Harley Books. pp. 67–118. part 2. Gelechiidae.
- Bland KP, Heckford RJ, Langmaid JR. 2002. Anacampsiniae. In: Emmet AM, Langmaid JR, editors. *The moths and Butterflies of Great Britain and Ireland*, vol. 4. Harley Books. pp. 204–220. part 2. Gelechiidae.
- CABI. 2019. *Fallopia japonica*. In: *Invasive Species Compendium*. Wallingford, UK: CAB International. Available at: <http://www.cabi.org/isc>. (Accessed 5 September 2019).
- Deschka G. 1982. *Bucculatrix pannonica* n. sp. (Lepidoptera, Bucculatricidae). *Zeitschrift Arbeitsgemeinschaft Österreichischer Entomologen* 34 (1/2):37–48.
- Diakonoff A. 1989. Revision of the Palaearctic Carposinidae with description of a new genus and new species (Lepidoptera: Pyraloidea). *Zoologische Verhandlungen* 251:1–155.
- Elsner G, Huemer P, Tokár Z. 1999. *Die palpenmotten (Lepidoptera, Gelechiidae) Mitteleuropas*. Bratislava. 208 pp.
- Emmet AM, editor. 1979. *A field guide to the Smaller British Lepidoptera*. British Entomological and Natural History Society. 271 pp.
- Emmet AM, editor. 1988. *A Field Guide to the Smaller British Lepidoptera*. British Entomological and Natural History Society. 288 pp.
- Emmet AM, editor. 1996. *The moths and butterflies of Great Britain and Ireland*. Harley Books. 452 pp.
- Emmet AM, Heath J, editors. 1991. *The Moths and Butterflies of Great Britain and Ireland*. Harley Books. 7(2). 353 pp.
- Emmet AM, Langmaid JR, editors. 2002. *The moths and butterflies of Great Britain and Ireland*. 4(1): Oecophoridae, Ethmiidae, Autostichidae, Blastobasidae, Batrachodidae, Agonoxenidae, Momphidae, Cosmopterigidae and Scythrididae. Harley Books. 326 pp.
- Falkovitsh MI. 2006. Host-plant relationships of the casebearers (Lepidoptera, Coleophoridae): Communication III. *Entomological Review* 86 (3):264–286.

- Fattorini S. 2013. Regional insect inventories require long time, extensive spatial sampling and good will. *PLoS One* 8 (4):e62118. <https://doi.org/10.1371/journal.pone.0062118>.
- Folmer O, Black M, Hoeh W, et al. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3:294–299.
- Gershenson ZS, Kozhevnikova VA. 2013. Lepidoptera, Plutellidae. In: Zerova MD, editor. *Fauna of Ukraine*, vol. 15. Kiev: Naukova Dumka. 135 pp. [in Russian].
- Hayes ML. 2019. *The biological control of weeds book: a New Zealand guide*. [Lincoln, N.Z.]: Manaaki Whenua, Landcare Research New Zealand. Available at: <https://www.landcareresearch.co.nz/publications/books/biocontrol-of-weeds-book>. (Accessed 5 September 2019).
- Heath J, editor. 1983. *The Moths and Butterflies of Great Britain and Ireland*, vol. 1. Harley Books. 344 pp.
- Heath J, editor. 1985. Emmet AM, editor. *The Moths and Butterflies of Great Britain and Ireland*, vol. 2. Harley Books. 460 pp.
- Hering M. 1957. Bestimmunstabellen der Blattminen von Europa: einschliesslich des Mittelmeerbeckens und der Kanarischen Inseln. *Junk, Gravenhage* 2:1–1185. 3:1–221.
- Huemer P, Hebert PDN. 2016. DNA barcode library for Lepidoptera from South Tyrol and Tyrol (Italy, Austria). *Impetus for integrative species discrimination in the 21st Century Gredleriana* 16 (X):41–164.
- Huemer P, Karsholt O. 1999. Gelechiidae I (Gelechiinae: Teleiodini, Gelechiini). In: *Microlepidoptera of Europe*, vol. 3. Stenstrup: Apollo Books. 354 pp.
- Huemer P, Mutanen M, Sefc KM, et al. 2014. Testing DNA barcode performance in 1000 species of European Lepidoptera: large geographic distances have small genetic impacts. *PLoS ONE* 9 (12):e115774.
- Huemer P, Wiesner M, Wiesmair B, et al. 2017. Schmetterlinge (Lepidoptera) des Altai-Gebirges (Südsibirien, Russland). Eindrücke einer internationalen Expedition im Spätsommer 2016. *Carinthia II* 207/127:527–564.
- Irisova NL, Bochkareva EN, Volynkin AV, et al. 2014. *Fauna and animal population. Nature annals of the Tigirekskii reserve, 2013. Book 11. "State reserve "Tigirekskii".* Barbaul. pp. 116–239.
- Ismagilov NN. 2018. Addition to the Katalog of animals of Bashkortostan on the order Lepidoptera (Insecta, Lepidoptera). In: Valuev VA, editor. *Material on flora and fauna of the Bashkortostan Republic*. Ufa: Publishing Centre BashSU. pp. 6–79.
- Izhevsky SS, Akhatov AK, Sinev SYu. 2011. Tomato mining moth already revealed in Russia. *Plant Protection and Quarantine* 3:40–44.
- Junnilainen J, Karsholt O, Nupponen K, et al. 2010. The gelechiid fauna of the southern Ural Mountains, part II: List of recorded species with taxonomic notes (Lepidoptera: Gelechiidae). *Zootaxa* 2367:1–68.
- Junnilainen J. 2001. Two new species of *Swammerdamia* from the southern Ural Mountains and southern Siberia (Lepidoptera: Yponomeutidae). *Entomologica Fennica* 12:112–118.
- Kenis M, Rabitsch W, Auger-Rozenberg MA, et al. 2007. How can alien species inventories and interception data help us prevent insect invasions? *Bulletin of Entomological Research* 97 (5):489–502. <https://doi.org/10.1017/S0007485307005184>.
- Kirichenko N, Triberti P, Kobayashi S, et al. 2018. Systematics of *Phyllocnistis* leaf-mining moths (Lepidoptera, Gracillariidae) feeding on dogwood (*Cornus* spp.) in Northeast Asia, with the description of three new species. *ZooKeys* 736:79–118. <https://doi.org/10.3897/zookeys.736.20739>.
- Kirichenko N, Triberti P, Lopez-Vaamonde C. 2019a. New species of leaf-mining *Phyllonorycter* (Lepidoptera Gracillariidae) from Siberia feeding on *Caragana* (Fabaceae). *Zookeys* 835:17–41. <https://doi.org/10.3897/zookeys.835.33166>.
- Kirichenko N, Triberti P, Mutanen M, et al. 2016. Systematics and biology of some species of *Micrurapteryx* Spuler (Lepidoptera, Gracillariidae) from the Holarctic Region, with re-description of *M. caraganella* (Hering) from Siberia. *Zookeys* 579:99–156. <https://doi.org/10.3897/zookeys.579.7166>.
- Kirichenko NI, Akulov EN, Ponomarenko MG, et al. 2019b. Molecular-genetic libraries of the leaf-mining micromoths (Lepidoptera: Gracillariidae) from Asian part of Russia and neighbouring territories. In: A.I. Kurentsov's Annual Memorial Meetings30; 2019. pp. 51–62. <https://sites.google.com/site/natakirisite/contact-us> [in Russian].
- Kirichenko NI, Akulov EN, Triberti P, et al. 2017a. New records of the leaf mining Gracillariid moths (Lepidoptera: Gracillariidae) from Asian part of Russia. *Far Eastern Entomologist* 346:1–12. <https://doi.org/10.25221/fee.346.1>.
- Kirichenko NI, Petko VM, Magnoux E, et al. 2017b. Diversity and distribution of leaf mining insects on Birches (*Betula* spp.) in Siberia. *Entomological Review* 97 (2): 183–198. <https://doi.org/10.1134/S0013873817020051>.
- Kirichenko NI, Triberti P, Akulov EN, et al. 2019c. Novel data on the taxonomic diversity, distribution and host plants of leafmining moths of the family Gracillariidae (Lepidoptera) in Siberia based on the DNA barcoding. *Entomological Review* 98 (3):600–631. <https://doi.org/10.1134/S0367144519030109> [in Russian].
- Klechkovsky YuE, Cherney LB, Vovkotrub ON. 2014. Tomato moth – new threat to agriculture. *Plant Protection and Quarantine* 4:36–39 [in Russian].
- Knyazev SA, Kirichenko NI, Baryshnikova SV, et al. 2018. The first records of the taxonomic diversity of leafmining micromoths, Gracillariidae (Insecta, Lepidoptera) in Omsk Province. *Eurasian Entomological Journal* 17:261–272 [in Russian].
- Koster JC, Sinev SYu. 2003. Momphidae, Batrachedridae, Stathmopodidae, Agonoxenidae, Cosmopterigidae, Chrysopoleiidae. In: *Microlepidoptera of Europe*, vol. 5. Stenstrup: Apollo Books. 387 pp.
- Kostjuk IYu, Budashkin YuI, Golovushkin MI. 1994. *Lepidoptera of the "Daurskii" reserve*. Kiev. 36 pp.
- Kozlov MV. 2008. Family Adelidae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. pp. 24–26 [in Russian].
- Kozlov MV. 2016. Fam. Adelidae. In: Beljaev EA, editor. *Annotated catalogue of the insects of Russian Far East*, vol. II. Lepidoptera. Vladivostok: Dalnauka. pp. 31–33 [in Russian].
- Kristensen NP, Scoble MJ, Karsholt O. 2007. Lepidoptera phylogeny and systematics: the state of inventorying moth and butterfly diversity. *Zootaxa* 1668:699–747.
- Kutenkova NN, Gorbach VV, Polevoi AV, et al. 2015. New data from Karelia for the Catalogue of Lepidoptera of Russia. *Eurasian Entomological Journal* 14 (4):299–308 [in Russian].
- Lepiforum: *Bestimmung von Schmetterlingen (Lepidoptera) und ihren Präimaginalstadien*. Available at: <http://www.lepiforum.de/>. (Accessed 20 July 2019).
- Lepidoptera. Part 1. In: Lerh PA, editor. 1997. *Key to the insects of Russian Far East*, vol. 5. Vladivostok: Dalnauka. 539 pp. [in Russian].
- Lepidoptera. Part 2. In: Lerh PA, editor. 1999. *Key to the insects of Russian Far East*, vol. 5. Vladivostok: Dalnauka. 671 pp. [in Russian].
- Lvovsky AL. 2006. Check-list of the broad-winged and flat moths (Lepidoptera: Oecophoridae, Chimabachidae, Amphibatidae, Depressariidae) of the fauna of Russia and adjacent countries. *Proceedings of the Zoological Institute Russian Academy of Sciences* 307:1–118 [in Russian].
- Lvovsky AL. 2008. Family Depressariidae. Family Chimabachidae. Family Oecophoridae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia* St.-Petersburg and Moscow: KMK Scientific Press. pp. 53–57. 65, 65–68. [in Russian].
- Lvovsky AL. 2016. Family Depressariidae. Family Chimabachidae. Family Oecophoridae. Lepidoptera. In: Beljaev EA, editor. *Annotated catalogue of the insects of Russian Far East*, vol. II. Vladivostok: Dalnauka. pp. 76–81. 88, 90–92. [in Russian].
- Maček V. 1999. *Hiponomoloska favna Slovenije = Hyponomologische Fauna Sloweniens*. Dela, slovenska Akademija Znanosti in Umetnosti. *Naravoslovne Vede* 37:1–385.
- Medvedev GS, editor. 1981. *The key to insects of the European part of the USSR. T. IV. Lepidoptera. Part 2*. Leningrad: "Nauka", Leningrad branch. 787 pp. [in Russian].
- Moth Dissection UK *Lepidoptera Dissection Group*; 2019. Available at: <https://mothdissection.co.uk/index.php>. (Accessed 20 July 2019).
- Nasu Y, Tamashima K, Shibao M, et al. 2010. Rediscovery of *Carposina niponensis* Walsingham and carposinids caught by synthetic sex pheromone trap for *C. sasakii* Matsumura in Japan (Lepidoptera: Carposinidae). *Japanese Journal of Applied Entomology and Zoology* 54:115–126.
- van Nieukerken EJ, Doorenweerd C, Hoare RJB, et al. 2016. Revised classification and catalogue of global Nepticulidae and Opostegidae (Lepidoptera, Nepticuloidea). *ZooKeys* 628:65–246. <https://doi.org/10.3897/zookeys.628.9799>.
- North American Moth Photographers Group in the Mississippi Entomological Museum at Mississippi State University. 2019. *Digital Guide to Moth Identification*. Available at: <https://mothphotographersgroup.msstate.edu/LatinIndex.shtml>. (Accessed 20 July 2019).
- Nupponen K. 2009. New records of Scythrididae from the Turanian region, with descriptions of seven new species (Lepidoptera: Scythrididae). *SHILAP Revista de Lepidopterología* 37 (147):341–362.
- Ponomarenko MG. 1997. Catalogue of the subfamily Dichomeriinae (Lepidoptera, Gelechiidae) of the Asia. *Far Eastern Entomologist* 50:1–67. URL: <http://www.biosoil.ru/Files/FEE/00000058.pdf>.
- Ponomarenko MG. 1999. Review of the family Carposinidae (Lepidoptera) from Russian Far East. *Far Eastern Entomologist* 69:1–12. URL: <http://www.biosoil.ru/Files/FEE/00000101.pdf>.
- Ponomarenko MG. 2008. Family Gelechiidae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. pp. 87–106 [in Russian].
- Ponomarenko MG. 2016. Fam. Tineidae. Fam. Argylethiidae. Fam. Ypsolophidae. Fam. Glyptopterigidae. Fam. Gelechiidae. Lepidoptera. In: Beljaev EA, editor. *Annotated catalogue of the insects of Russian Far East*, vol. II. Vladivostok: Dalnauka. pp. 36–43. 64–69, 72–73, 115–139. [in Russian].
- Ponomarenko MG, Omelko MM. 2003. Review of the genus *Acanthophila* Heinemann, 1870 (Lepidoptera, Gelechiidae). *Far Eastern Entomologist* 127:1–24. URL: <http://www.biosoil.ru/Files/FEE/00000175.pdf>.
- Razowski J, Kumata T. 1985. Typenkatalog der von S. Matsumura beschriebenen Lepidopteren. *Neue Entomologische Nachrichten* 17:1–28.
- Robinson GS. 1976. The preparation of slides of Lepidoptera genitalia with special reference to the Microlepidoptera. *Entomologist's Gazette* 27:127–132.
- Rosellkhonadzor: *Normativnye dokumenty*. Available at: <https://www.fsvps.ru/fsvps/laws>. (Accessed 9 September 2019).
- Sattler K. 1980. *Telediodes wagae* (Nowick, 1860) New to the British List (Lepidoptera, Gelechiidae). *Entomologist's Gazette* 31:235–245.
- Savela M. 2001. *Lepidoptera and Some Other Life Forms*. Available at: <http://www.nic.funet.fi/>. (Accessed 13 May 2019).
- Savotikov YuF, Smetnik AI. 1995. Reference book on the pests, diseases and weeds of quarantine significance for the territory of the Russian Federation. *Arnika, Nizhny Novgorod*:230 [in Russian].
- Sinev SYu, editor. 2008a. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. 424 pp. [in Russian].
- Sinev SYu. 2008b. Family Nepticulidae. Family Yponomeutidae. Family Ypsolophidae. Family Glyptopterigidae. Family Elachistidae. Family Momphidae. Family

- Cosmopterigidae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. pp. 19–23. 45–48, 51, 58–61, 82–83, 85–87. [in Russian].
- Sinev SYu. 2016. Fam. Psychidae. Fam. Elachistidae. Fam. Cosmopterigidae. Lepidoptera. In: Beljaev EA, editor. *Annotated catalogue of the insects of Russian Far East*, vol. II. Vladivostok: Dalnauka. pp. 43–46. 82–84, 110–114. [in Russian].
- Sinev SYu, Lovtsova YuA. 2008. Family Psychidae. In: Sinev SYu, editor. *Catalogue of the Lepidoptera of Russia*. St.-Petersburg and Moscow: KMK Scientific Press. pp. 32–36 [in Russian].
- Sohn JC. 2009. Two species of Yponomeutidae (Lepidoptera) new to Korea with the first description of female of *Cedestis exiguata* Moriuti. *Tinea* 20 (5):294–298.
- Stainton HT. 1851. *A Supplementary Catalogue of the British Tineidæ and Pterophoridaæ*. John van Voorst. 28 pp.
- Suckling DM, Conlong DE, Carpenter JE, et al. 2017. Global range expansion of pest Lepidoptera requires socially acceptable solutions. *Biological Invasions* 19 (4): 1107–1119. <https://doi.org/10.1007/s10530-016-1325-9>.
- Svenska. 2019. Svenska fjärilar Lepidoptera. Available at: http://www2.nrm.se/en/svenska_fjarilar/svenska_fjarilar.html. (Accessed 20 July 2019).
- Traugott-Olsen E, Nielsen ES. 1977. The Elachistidae (Lepidoptera) of Fennoscandia and Denmark. *Fauna Entomologica Scandinavica*, Klampenborg 6:1–299.
- van Roosmalen JAM, Doorenweerd C. 2015. *Coleophora gryphipennella* (Hübner, 1796) (Lepidoptera, Coleophoridae) on *Fragaria vesca* L. (Rosaceae), a novel host, in the coastal dunes of The Netherlands. *Nota Lepidopterologica* 38 (2):147–155. <https://doi.org/10.3897/nl.38.6289>.
- Winston RL, editor. 2014. *Biological control of weeds: a world catalogue of agents and their target weeds*. 5th edition. Morgantown, West Virginia: USDA Forest Service, Forest Health Technology Enterprise Team. FHTET-2014-04. 838 pp.
- Zhimerikin VN, Mironova MK. 2012. South American tomato moth – threat for tomato production. *Plant Protection and Quarantine* 11:32–34 [in Russian].