

A brief review of the genus *Mantura* Stephens, 1831 (Coleoptera: Chrysomelidae: Galerucinae: Alticini) of Russia and some adjacent territories

© M.Ye. Sergeev

Federal Scientific Center of the East Asia Terrestrial Biodiversity of the Far Eastern Branch of the Russian Academy of Sciences, 100 let Vladivostoku Av., 159, Vladivostok 690022 Russia. E-mail: eksgauster@inbox.ru

Abstract. Seven species of the genus *Mantura* Stephens, 1831 are reviewed. In total, six species are known in Russia. *Mantura rustica* (Linnaeus, 1767), *M. chrysanthemii* (Koch, 1803), *M. pallidicornis* (Waltl, 1839) and *M. cylindrica* Miller, 1881 are distributed in the European part of the country. One European species, *Mantura obtusata* (Gyllenhal, 1813) needs verification for distribution in Russia. Two species, *M. japonica* Jacoby, 1885 and *M. clavareauii* Heikertinger, 1912 are recorded for the fauna of Russia for the first time. Both species were collected in the Russian Far East, the first one in Primorsky Region, the latter on Sakhalin. The genus *Mantura* is newly recorded from Sakhalin Island. Key to species of *Mantura* of the Russian fauna is given.

Key words: *Mantura*, faunistics, new records, identification key, Palaearctic, Russia, Sakhalin.

Краткий обзор рода *Mantura* Stephens, 1831 (Coleoptera: Chrysomelidae: Galerucinae: Alticini) фауны России и некоторых прилегающих территорий

© М.Е. Сергеев

Федеральный научный центр биоразнообразия наземной биоты Восточной Азии ДВО РАН, пр. 100 лет Владивостоку, 159, Владивосток 690022 Россия. E-mail: eksgauster@inbox.ru

Резюме. Приведен обзор семи видов рода *Mantura* Stephens, 1831 фауны России и сопредельных территорий. *Mantura rustica* (Linnaeus, 1767), *M. chrysanthemii* (Koch, 1803), *M. pallidicornis* (Waltl, 1839) и *M. cylindrica* Miller, 1881 распространены в европейской части страны. Для одного европейского вида, *M. obtusata* (Gyllenhal, 1813), требуется верификация материала для подтверждения его распространения в России. Два таксона, *M. japonica* Jacoby, 1885 и *M. clavareauii* Heikertinger, 1912, впервые указаны для фауны России. Оба вида были найдены на Дальнем Востоке, первый в Приморском крае, второй на Сахалине (род *Mantura* впервые указывается для фауны острова Сахалин). Дана определительная таблица видов фауны России.

Ключевые слова: *Mantura*, фаунистика, новые указания, определительная таблица, Палеарктика, Россия, Сахалин.

Introduction

The genus *Mantura* Stephens, 1831 is divided into two subgenera (*Mantura* s. str. and *Stenomantura* Heikertinger, 1909), with 20 species in the world, 15 of them are distributed in the Palaearctic. Several taxa occur in the North America (Canada, USA), Africa and Southeast Asia [Lopatin, 1977, 2010; LeSage, 1991; Kimoto, Takizawa, 1994; Konstantinov, Vandenberg, 1996; Riley et al., 2003; Gruev, Döberl, 2005; Medvedev, 2009; Döberl, 2010; Warchałowski, 1998, 2010]. Five species of *Mantura* were known so far in the fauna of Russia: *M. rustica* (Linnaeus, 1767), *M. chrysanthemii* (Koch, 1803), *M. obtusata* (Gyllenhal, 1813), *M. pallidicornis* (Waltl, 1839) and *M. cylindrica* Miller, 1881 [Gressitt, Kimoto, 1963; Medvedev, Shapiro, 1965; Medvedev, 1982, 1992; Kimoto, Takizawa, 1994; Warchałowski, 1998, 2010; Bieńkowski, 2004; Lopatin et al., 2004; Gruev, Döberl, 2005; Lopatin, Nesterova, 2005; Döberl, 2010; Dolgin, Bieńkowski, 2011; Sergeev, 2017, 2018]. Species of *Mantura* inhabit mesophytic biotopes and trophically are associated with Polygonaceae, Asteraceae, Cistaceae [Medvedev, Roginskaya, 1988; Jolivet, Hawkeswood, 1995;

Warchałowski, 1998; Baviera, Biondi, 2015]. Adults feed on skeletonize leaves, while larvae develop in the mines inside of leaves, possibly also in the roots of a plant in the soil [Zaitsev, Medvedev, 2009].

In this paper two additional species, *M. japonica* Jacoby, 1885 and *M. clavareauii* Heikertinger, 1912 are recorded for Russia. The distribution of *M. obtusata* (Gyllenhal, 1813) and *M. pallidicornis* (Waltl, 1839) is clarified.

Material and methods

This work is based on the material collected by the author in 2015–2019 in Primorskiy Region and Sakhalin Island, and deposited in the collection of Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far Eastern Branch of the Russian Academy of Sciences (EATB, Vladivostok, Russia). Materials from the following collections also have been studied:

MNKNU – V.N. Karazin State Museum of Nature Kharkov National University (Kharkov, Ukraine);

IZS – I.I. Schmalhausen Institute of Zoology of the National Academy of Sciences of Ukraine (Kiev, Ukraine);

NSU – N.V. Gogol Nezhyn State University (Nezhyn, Ukraine);

SNHM – State Natural History Museum of the National Academy of Sciences of Ukraine (Lvov, Ukraine);

ZIN – Zoological Institute of the Russian Academy of Sciences (St Petersburg, Russia).

Photographs were taken with the stereomicroscope Olympus SZX16 and digital camera Olympus DP74 and stacked using Helicon Focus software. The final illustrations were post-processed for contrast and brightness using Adobe Photoshop® software.

The classification of Chrysomelidae based on Riley et al. [2003] is used.

Subfamily Galerucinae Latreille, 1802

Tribe Alticini Newman, 1835

Genus *Mantura* Stephens, 1831

Mantura Stephens, 1831: 322. Type species: *Chrysomela rustica* Linnaeus, 1767, by subsequent designation [Westwood, 1840].

Cardiapus Curtis, 1833: pl. 435. Type species: *Cardiapus mathewsii* Curtis, 1833, by monotypy.

Balanomorpha Chevrolat in Dejean, 1836: 393. Type species: *Galleruca semiaenea* Fabricius, 1792 (= *Chrysomela rustica* Linnaeus, 1767), by subsequent designation of Heikertinger [1951]. Junior subjective synonym of *Mantura* according to Weise [1886].

Mantura (Mantura) chrysanthemii (Koch, 1803)

(Figs 1–3)

Haltica chrysanthemii Koch, 1803: 45 (type locality: Palatinate, Germany).

Mantura chrysanthemii: Zhyvotovskaya, 1957: 52; Medvedev, Shapiro, 1965: 454; Bieńkowski, 2004: 92; Gus'kova, 2010: 217; Dolgin, Bieńkowski, 2011: 232; Ruchin, 2011: 167; Sergeev, Sheshurak, 2014: 23; Sergeev, 2018: 85.

Material. Ukraine: 1 ex., (IZS), Donetsk Region, “Kamennye Mogily” Nature Reserve, 2.06.1970 (V.G. Dolin); 1 ex., (EATB), same locality, 26.05.1998 (E.S. Ivanova); 1 ex., (EATB), same locality, 19.06.2004 (M.Ye. Sergeev); 1 ex., (EATB), same locality, 15.05.2006 (E.Yu. Savchenko); 1 ex., (EATB), same locality, 15.06.2006 (E.Yu. Savchenko); 2 ex., (EATB), Slavyansk District, Slavyansk env., 22.04.2002 (V.Yu. Nazarenko); 1 ex., (NSU), Chernigov Region, Sednev env., floodplain of the Snov River, 10–13.06.1991 (P.N. Sheshurak); 2 ex., (EATB), Kiev, coast of Telbin Lake, 1.09.2000 (V.Yu. Nazarenko); 2 ex., (EATB), same locality, 20.06.2002 (V.Yu. Nazarenko); 1 ex., (SNHM), Lvov env., 11.06. (clear date and collector unknown).

Distribution. North-West Africa, Europe. Russia: European part (Leningrad, Moscow, Vladimir, Tula and Penza regions, Komi Republic, Mordovian Republic, Udmurt Republic), Ural (Perm Region, Bashkir Republic) [Zhyvotovskaya, 1957; Bieńkowski, 2004; Gus'kova, 2010; Warchałowski, 2010; Ruchin, 2011; Dolgin, Bieńkowski, 2011; Bieńkowski, Orlova-Bieńkovskaya, 2012; Sergeev, Sheshurak, 2014; Sergeev, 2018]. Introduced in North America (Canada, USA) [LeSage, 1991; Riley et al., 2003].

Mantura (Mantura) clavareai Heikertinger, 1912

(Figs 4–6)

Mantura (Mantura) clavareai Heikertinger, 1912: 45 (syntypes from Japan: “Jesso” (Hokkaido), “Kioto” (Kyoto, Honshu)).

Mantura (Mantura) clavareai: Kimoto, Takizawa, 1994: 248; Warchałowski, 2010: 910; Cho, An, 2020: 18.

Material. Russia: 1♂, (EATB), Sakhalin, Nevelsk District, Yasnomorskoe env., floodplain of Yasnomorka River, 19.07.2019 (M.Ye. Sergeev).

Distribution. Russia (new record): Sakhalin. South Korea, Japan: Hokkaido, Honshu, Kyushu, Shikoku [Kimoto, Takizawa, 1994; Warchałowski, 2010; Cho, An, 2020].

Mantura (Mantura) japonica Jacoby, 1885

(Figs 7, 8)

Mantura (Mantura) japonica Jacoby, 1885: 720 (type locality: “Hakodate” (Hokkaido, Japan)).

Mantura japonica: Kimoto, Takizawa, 1994: 248; Warchałowski, 2010: 910.

Material. Russia: 1♀, (EATB), Primorskiy Region, Khasansky District, near Barabash, meadow in floodplain of Barabashevka River, 8.06.2019 (M.Ye. Sergeev).

Distribution. Russia (new record): Primorskiy Region. Japan: Hokkaido [Kimoto, Takizawa, 1994; Warchałowski, 2010].

Mantura (Mantura) obtusata (Gyllenhal, 1813)

Haltica obtusata Gyllenhal, 1813: 579 (type locality: Sweden, according to Wanntorp [2008: 100]).

Haltica ambigua Kutschera, 1862: 52. (type locality: Aachen, Germany). Junior subjective synonym of *Haltica obtusata* according to Wanntorp [2008].

Mantura obtusata: Wanntorp, 2008: 129; Döberl, 2010: 536.

Notes. Döberl [2010] did not list this species from Russia. However, this species was indicated for the north-west of the European part of Russia (Pskov, Yaroslavl' and Moscow regions) [Bieńkowski, 2004; Bieńkowski, Orlova-Bieńkovskaya, 2012; Vlasov, Rusinov, 2017]. We have not studied this material, but we can assume that *M. obtusata* occurs in the north-west of the European part of Russia on the basis of distribution of this species in western and northern Europe.

Distribution. Western and Northern Europe, in Eastern Europe it is known from Poland (Lower Silesia); in Southern Europe it is known in the northern part of the Balkan Peninsula (Slovenia) [Warchałowski, 1998; Bieńkowski, 2004; Gruev, Döberl, 2005; Borowiec et al., 2011].

Mantura (Mantura) pallidicornis (Waltl, 1839)

(Figs 9–11)


Haltica pallidicornis Waltl, 1839: 225 (type locality: “Passau”, Germany).

Mantura pallidicornis: Wanntorp, 2008: 100; Döberl, 2010: 536.

Mantura obtusata: Medvedev, Shapiro, 1965: 454; Warchałowski, 1998: 178; Bieńkowski, 2004: 92; Sergeev, 2018: 86.

Material. Ukraine: 1 ex., (MNKNU), Kharkov Region, Petrovsky District, Protopyovka, 21.06.1952 (D.S. Shapiro); 2 ex., (NSU), Chernigov Region, Koropsky District, Obolonie env., Desna River valley, 17.06.1992 (P.N. Sheshurak).

Notes. Before the work of Wanntorp [2008] *Mantura pallidicornis* was unknown from Eastern Europe, including Russia and Ukraine. Wanntorp [2008] examined genitalia (aedeagus) of *M. obtusata* and clearly separated this species from the close *M. pallidicornis*. According to his data, the distribution of *M. obtusata* is limited by countries



Figs 1–14. *Mantura* species, habitus and details of structure.

1–3 – *M. chrysanthemi*, male; 4–6 – *M. clavareai*, male; 7–8 – *M. japonica*, female; 9–11 – *M. pallidicornis*, male; 12–14 – *M. rustica*, male. 1, 4, 7, 9, 12 – habitus; 2, 5, 10, 12 – aedeagus, ventral view; 3, 6, 11, 14 – the same, lateral view; 8 – spermatheca.

Рис. 1–14. Виды рода *Mantura*, габитус и детали строения.

1–3 – *M. chrysanthemi*, самец; 4–6 – *M. clavareai*, самец; 7–8 – *M. japonica*, самка; 9–11 – *M. pallidicornis*, самец; 12–14 – *M. rustica*, самец. 1, 4, 7, 9, 12 – габитус; 2, 5, 10, 12 – эдеагус, вентрально; 3, 6, 11, 14 – то же, латерально; 8 – сперматека.

of Western and Northern Europe to the North Balkans on south, while *M. pallidicornis* is widely distributed in Europe [Döberl, 2010].

Distribution. Europe (Northern, Central, Southern, Eastern, including Ukraine and south of the European part of Russia) [Döberl, 2010].

Mantura (Mantura) rustica (Linnaeus, 1767)
(Figs 12–14)

Chrysomela rustica Linnaeus, 1767: 595 (type locality: “ad Hammerby”, Sweden).

Mantura rustica: Medvedev, Shapiro, 1965: 454; Medvedev, 1992: 594; Kubisz et al., 1997: 263; Warchałowski, 1998: 179–180; Bieńkowski, 2004: 92; Gus’kova, 2010: 217; Sergeev, 2017: 200; Sergeev, 2018: 86; Cho, An, 2020: 18.

Material. Russia: 1 ex., (ZIN), Primorskiy Region, Anuchino District, Vinogradovka env., 28.06.1929 (Kirichenko); 1 ex., (EATB), Primorskiy Region, Kirovsky District, Pavlo-Fedorovka env., Khanka Nature Reserve, 12.07.2016 (M.Ye. Sergeev); 1 ex. (EATB), Novosibirsk Region, Kolyvan’ District, Tropino env., birch forest, 21.07.1978 (V.N. Kuznetsov). Ukraine: 2 ex. (EATB), Donetsk, 2–31.05.1999 (E.V. Prokopenko); 2 ex. (EATB), Lugansk Region, near Dyakovo, 12.04.2000 (S.V. Kononov); 1 ex. (EATB), Stanitsa Luganskaya District, Lugansk Nature Reserve, floodplain of the Seversky Donets River, 18.06.2002 (A.G. Maltseva); 2 ex., (EATB), same locality, 12.06.2003 (A.G. Maltseva); 2 ex., (EATB), Dnepropetrovsk Region, Sinel’niko District, Raevka, 15.06.1984 (A.M. Sumarokov).

Distribution. The widespread Palearctic species. Western and Eastern Europe (including Ukraine). Russia: European part (Moscow, Oryol, Volgograd regions, Kalmyk Republic, Udmurt Republic, Crimea), Ural, Western Siberia (Novosibirsk Region, Altai), south of Eastern Siberia (Krasnoyarsk and Irkutsk regions, Yakutsk Republic), the Far East (Amur and Primorskiy regions), Turkey, Kazakhstan, Kyrgyzstan, Tajikistan, Afghanistan, Mongolia, China, South Korea [Lopatin, 1977, 2010; Medvedev, 1992; Kubisz et al., 1997; Warchałowski, 1998, 2010; Mosyakin, Popov, 1999; Bieńkowski, 2004; Gus’kova, 2010; Döberl, 2010; Bieńkowski, Orlova-Bieńkovskaya, 2012; Sergeev, 2017, 2018; Cho, An, 2020].

Mantura (Stenomantura) cylindrica Miller, 1881

Mantura cylindrica Miller, 1881: 2 (type locality: “Dalmatia” (Croatia)).

Mantura cylindrica: Medvedev, Shapiro, 1965: 454; Warchałowski, 1998: 181; Gruev, Döberl, 2005: 112; Döberl, 2010: 537.

Distribution. Europe (Italy (Sicily), Croatia, Bulgaria, Greece), Asian Turkey, Syria, Azerbaijan, Russia (Dagestan) [Medvedev, Shapiro, 1965; Warchałowski, 1998; Gruev, Döberl, 2005; Döberl, 1999, 2010; Baviera, Biondi, 2015].

Key to the *Mantura* species of Russia

(based on Medvedev, Shapiro [1965],
Kimoto, Takizawa [1994],
Bieńkowski [2004], Warchałowski [2010]
with modifications)

1. Pronotum anterad not narrowed, posterolaterally with distinct longitudinal short groove. Elytra posterad longitudinally acute. (Subgenus *Mantura*) 2
- Pronotum anterad strongly narrowed and projected over head; posterolaterally without distinct longitudinal short groove. Elytra posterad almost rounded.

- (Subgenus *Stenomantura*). – Body length 1.8–2 mm *M. (S.) cylindrica*
2. Elytra unicolorous 3
- Elytra bicolorous (Fig. 12). – Body length 2–2.8 mm *M. (M.) rustica*
3. Body dorsally shining 4
- Body dorsally matte 5
4. Body dorsally dark metallic green (Fig. 4). Pronotum dorsally distinctly punctate and finely granulate. Legs reddish brown, often hind femora bronzy. – Body length 2.5–3 mm. Aedeagus (Fig. 5, 6) *M. (M.) clavareau*
- Body dorsally pitchy-brown or brownish-red, mostly with bronze reflection, or dark bronze (Fig. 1). Legs rust-reddish or dark amber, hind femora and apical part of antennae darkened. – Body length 1.8–2.7 mm. Aedeagus (Fig. 2, 3) *M. (M.) chrysanthem*
5. Body dorsally black with obscure greenish or bluish tint 6
- Body dorsally dark blue (Fig. 7). – Spermatheca (Fig. 8). Body length 1.8–2.1 mm *M. (M.) japonica*
6. Aedeagus ventrally without distinct rills (Figs 10, 11). Body length 1.8–2.6 mm (Fig. 9) ... *M. (M.) pallidicornis*
- Aedeagus ventrally with distinct rills. Body length 1.8–2 mm *M. (M.) obtusata*

Discussion

The genus *Mantura* is represented by seven species in the fauna of Russia, including two species (*Mantura japonica* and *M. clavareau*) new for the fauna of the country; the genus is also recorded for the fauna of Sakhalin Island for the first time.

Species of this genus are not numerous in nature and usually are presented by single specimens or small series in collection materials. Therefore, new records are valuable additions to the current knowledge of the distribution of *Mantura* within the fauna of Russia and adjacent territories. Endemic species are absent in the fauna of the country. The mentioned species belong to the four biogeographical groups: Transpalearctic – *M. rustica*; Western Palearctic – *M. obtusata*, *M. pallidicornis* and *M. chrysanthem*, the latter was introduced in North America; Mediterranean – *M. cylindrica*; Eastern Palearctic – *M. japonica* and *M. clavareau*.

One species, *M. (s. str.) mathewsii* Curtis, 1833 is distributed in Western, Central, Southern and Eastern Europe (including Ukrainian Carpathians), Azerbaijan (Absheron Peninsula), Asian Turkey (Denizli, Isparta provinces) [Medvedev, Shapiro, 1965; Warchałowski, 1998; Lopatin et al., 2004; Gruev, Döberl, 2005; Döberl, 2010; Ekiz et al., 2013] and can be found in the North Caucasus (Dagestan Republic or Krasnodar Region of Russia).

Acknowledgements

The author thanks E.A. Belyaev, M.G. Ponomarenko, V.M. Loktionov, M.Yu. Proshchalykin and A.S. Lelej (EATB) for their help during collection trip and valuable suggestions on this work, Jan Bezděk (Mendel University, Brno, Czech Republic) and A.G. Moseyko (Zoological Institute of the

Russian Academy of Sciences, St Petersburg, Russia) for valuable comments and corrections. The author is also grateful to E.S. Ivanova, A.G. Maltseva, E.V. Prokopenko, E.Yu. Savchenko (Donetsk, Ukraine), S.V. Konovalova (Dyakovo, Ukraine), A.N. Drovalenko (Kharkov, Ukraine), A.M. Sumarokov (Raevka, Ukraine), V.Yu. Nazarenko (Kiev, Ukraine), P.N. Sheshurak (Nezhin, Ukraina), T.P. Yanitsky (L'vov, Ukraine), A.G. Moseyko for the loan material.

References

- Baviera C., Biondi M. 2015. The Alticini (Coleoptera: Chrysomelidae, Galerucinae) of Sicily: recent records and updated checklist. *Atti della Accademia Peloritana dei Pericolanti*. 93(2): 1–50. DOI: 10.1478/AAPP932A2
- Bieńkowski A.O. 2004. Leaf-beetles (Coleoptera: Chrysomelidae) of the Eastern Europe. New key to subfamilies, genera, and species. Moscow: Mikron-print. 278 p.
- Bieńkowski A.O., Orlova-Berikowskaya M.Ya. 2012. Catalogue of localities of leaf beetles (Chrysomelidae) of Russia and adjacent regions. Version 3.11.2012. In: Beetles (Coleoptera) and coleopterists. Available at: <https://www.zin.ru/animalia/coleoptera/rus/benkat11.htm> (in Russian) (accessed 18 September 2020).
- Borowiec L., Scibior R., Kubisz D. 2011. Critical check-list of the Polish Chrysomeloidea, excluding Cerambycidae (Coleoptera: Phytophaga). *Genus*. 22(4): 579–608.
- Chevrolat L.A.A. 1836. [New taxa]. In: Dejean P.F.A.M. Catalogue des coléoptères de la collection de M. le Comte Dejean, Deuxième édition. Livraison 5. Paris: Méquignon-Marvis Père et Fils: 361–442.
- Cho H.-W., An S.L. 2020. An annotated checklist of leaf beetles (Coleoptera: Chrysomelidae) of Korea, with comments and new records. *Far Eastern Entomologist*. 404: 1–36. DOI: 10.25221/fee.404.1
- Curtis J. 1833. British entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland: containing coloured figures from nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. Vol. X. London: J. Curtis: 434–481.
- Döberl M. 1999. Zur Alticinenfauna von Azerbaijan. *Mitteilungen des entomologischen Vereins*. 34: 129–130.
- Döberl M. 2010. Subfamily Alticinae Newman, 1835. In: Catalogue of Palaearctic Coleoptera. Vol. 6. Chrysomeloidea. (I. Löbl, A. Smetana eds). Senstrup: Apollo Books: 491–563.
- Dolgin M.M., Bieńkowski A.O. 2011. Fauna evropeyskogo severo-vostoka Rossii. T. 8. Vyp. 3. Zhuki-listoedy (Coleoptera, Chrysomelidae) [Fauna of the European Northeast of Russia. Vol. 8. Iss. 3. Leaf beetles (Coleoptera, Chrysomelidae)]. St Petersburg: Nauka. 291 p. (in Russian).
- Ekiz A.N., Şen I., Aslan E.G., Gök A. 2013. Check list of leaf beetles (Coleoptera: Chrysomelidae) of Turkey, excluding Bruchinae. *Journal of Natural History*. 47(33–34): 2213–2287.
- Gressitt J.L., Kimoto S. 1963. The Chrysomelidae (Coleoptera) of China and Korea. Part 2. *Pacific Insect Monograph*. 1B: 743–893.
- Gruev B., Döberl M. 2005. General distribution of the flea beetles in the Palaearctic subregion (Coleoptera, Chrysomelidae: Alticinae). Supplement. Sofia–Moscow: Pensoft. 240 p.
- Gus'kova E. 2010. The leaf beetles (Coleoptera, Chrysomelidae) of the South Urals. *Entomofauna. Zeitschrift für Entomologie*, 31(14): 169–228.
- Gyllenhal L. 1813. Insecta Suecica descripta. Classis I. Coleoptera sive Eleutherata. Tom 1, Pars III. Scaris: Leverents: [4] + 730 + [2] p.
- Heikertinger F. 1912. Diagnosen neuer paläarktischer Halticinen. *Verhandlungen der Kaiserlich-Königlichen Zoologisch-Botanischen Gesellschaft in Wien*. 62: 44–48.
- Heikertinger F. 1951. Bestimmungstabellen der paläarktischen Arten der Gattungen *Podagrica* Foudr., *Mantura* Steph. und *Chaetocnema* Steph. *Koleopterologische Rundschau*. 32: 1–84.
- Jacoby M. 1885. Description of the phytophagous Coleoptera of Japan obtained by Mr. George Lewis during his second journey, from February 1880 to September 1881. – Part II. Halticini and Galerucinae. *Proceedings of the Scientific Meetings of the Zoological Society of London*. 1885: 719–755, pl. XLVI.
- Jolivet P., Hawkeswood T.L. 1995. Host-plants of the Chrysomelidae of the world. An essay about the relationships between the leaf-beetles and their food-plants. Leiden: Backhuys Publishers. 281 p.
- Kimoto Sh., Takizawa H. 1994. Leaf beetles (Chrysomelidae) of Japan. Tokyo: Tokai University Press. 539 p.
- Koch J.D.W. 1803. I. Monographie der von den Verfassern in dem Departemente vom Donnersberge, und den angrenzenden Gegenden der Departemente von der Saar, und von Rhein und Mosel einheimisch beobachteten Flohkäfer (Haltica). In: Entomologische Hefte enthaltend Beiträge zur weitem Kenntniss und Aufklärung der Insektengeschichte. Eine Vorarbeit zu einer künftigen Fauna des Departements vom Donnersberge. Zweites Heft. Frankfurt am Main: Im Verlage bei Friedrich Efslinger: 1–130.
- Konstantinov A.S., Vandenberg N.J. 1996. Handbook of Palearctic Flea Beetles (Coleoptera, Chrysomelidae: Alticinae). Contribution on Entomology, International, 1(3). Gainesville: Associated Publishers. 439 p.
- Kubisz D., Mazur M., Pawlowski J. 1997. Chrzaszczce Miodoborow (Zachodnia Ukraina) Czesc II. Aktualny stan poznania (Insecta: Coleoptera). *Studia Ośrodka Dokumentacji Fizjograficznej*. 25: 217–294.
- Kutschera F. 1862. Beiträge zur Kenntnis europäischer Halticinen. *Wiener Entomologische Monatsschrift*. 6: 47–55, 97–109, 215–228.
- LeSage L. 1991. Chrysomelidae. In: Checklist of beetles of Canada and Alaska. Ottawa: Biosystematic Research Centre, Research Branch Agriculture Canada: 301–323.
- Linnaeus C. 1767. Systema Nature per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus differentibus, synonymis, locis. Editio duodecima, reformata. Tom. 1. Pars II. Homiae: Laurentii Salvii: [2] + 533–1327 + [37] p.
- Lopatin I.K. 1977. Opredeliteli po faune SSSR. Zhuki-listoedy (Chrysomelidae) central'noy Azii i Kazakhstana [Key to the fauna of the USSR. Leaf beetles (Chrysomelidae) of Central Asia and Kazakhstan]. Leningrad: Nauka. 270 p. (in Russian).
- Lopatin I.K. 2010. Zhuki-listoedy (Insecta, Coleoptera, Chrysomelidae) Central'noy Azii [Leaf beetles (Insecta, Coleoptera, Chrysomelidae) of Central Asia]. Minsk: Belarusian State University. 511 p. (in Russian).
- Lopatin I.K., Aleksandrovich O.R., Konstantinov A.S. 2004. Check list of leaf beetles (Coleoptera, Chrysomelidae) of the Eastern Europe and Northern Asia. Olsztyn: Mantis. 343 p.
- Lopatin I.K., Nesterova O.L. 2005. Nasekomye Belarusi: Zhuki-listoedy (Coleoptera, Chrysomelidae) [Insects of Belarus: Leaf beetles (Coleoptera, Chrysomelidae)]. Minsk: Technoprint. 294 p. (in Russian).
- Medvedev L.N. 1982. Zhuki-listoedy MNR: Opredelitel' [Leaf beetles of the Mongolian People's Republic: Identification keys]. Moscow: Nauka. 301 p. (in Russian).
- Medvedev L.N. 1992. Opredelitel' nasekomykh Dal'nego Vostoka SSSR. T. III. Zhestkokrylye, ili zhuki. Ch. 2. Chrysomelidae – Zhuki-listoedy [Key to the insects of Far East of the USSR. Vol. III. Coleoptera. Part 2. Chrysomelidae]. St Petersburg: Nauka: 533–602 (in Russian).
- Medvedev L.N. 2009. Alticinae of Indochina. Moscow: KMK Scientific Press Ltd. 224 p.
- Medvedev L.N., Roginskaya E.Ya. 1988. Katalog kormovykh rasteniy zhukov-listoedov SSSR [Catalog of host plants of leaf beetles of the USSR]. Moscow: PEM VNIIS Gosstroya of the USSR. 192 p. (in Russian).
- Medvedev L.N., Shapiro D.S. 1965. Opredelitel' nasekomykh evropeyskoy chasti SSSR. T. 2. Vyp. 89. Chrysomelidae – zhuki-listoedy [Key to the fauna of the European part of the USSR. Vol. 2. Iss. 89. Chrysomelidae]. Moscow – Leningrad: Nauka: 419–474 (in Russian).
- Miller L. 1881. Bericht über eine im Frühling 1879 nach Dalmatien unternommene coleopterologische Reise. *Verhandlungen der Kaiserlich-Königlichen Zoologischen-Botanischen Gesellschaft in Wien*. 1880. 30: 1–8.
- Mosyakin S.A., Popov V.N. 1999. Ecological and faunistic structure of the fauna of leaf beetles (Coleoptera, Chrysomelidae) of Crimea. In: Prilozhenie k nauchno-prakticheskomu diskussionno-analiticheskomu sborniku "Voprosy razvitiya Kryma" [Supplement to the scientific-practical discussion-analytical collection "Problems of the development of Crimea"]. Simferopol: Sonat: 30–40 (in Russian).
- Riley E.G., Clark S.M., Seeno T.N. 2003. Catalog of the leaf beetles of America north of Mexico (Coleoptera: Megalopodidae, Orsodacnidae and Chrysomelidae, excluding Bruchinae). Sacramento, CA: The Coleopterists Society. 290 p.
- Ruchin A.B. 2011. The first additional materials to the entomofauna of the Mordovian State Nature Reserve. *Trudy Mordovskogo gosudarstvennogo prirodnogo zapovednika imeni P.G. Smidovicha*. 9: 150–182 (in Russian).
- Sergeev M.E. 2017. Materials for the fauna of leaf beetles (Coleoptera: Chrysomelidae: Megalopodidae) of the Khanka Reserve. *Trudy Mordovskogo gosudarstvennogo prirodnogo zapovednika imeni P.G. Smidovicha*. 19: 189–205. (in Russian).
- Sergeev M.E. 2018. Leaf beetles (Coleoptera: Megalopodidae, Chrysomelidae, Orsodacnidae) in southeastern Ukraine. In: *Trudy*

- Russkogo entomologicheskogo obshchestva. T. 89 [Proceedings of the Russian Entomological Society. Vol. 89]. St Petersburg: Zoological Institute of the Russian Academy of Sciences: 1–121 p. (in Russian).
- Sergeev M.E., Scheschurak P.N. 2014. Leaf beetles (Coleoptera, Chrysomelidae) of the Forest zone of Ukraine. *Ukrainian Entomological Journal*. 2(9): 16–29 (in Russian).
- Stephens J.F. 1831. *In: Illustration of British entomology; or, a synopsis of indigenous insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as practicable. Mandibulata. Volume IV.* London: Baldwin & Cradock: 1–366.
- Vlasov D.V., Rusinov A.A. 2017. Fauna of leaf beetles (Coleoptera, Chrysomelidae) of the Pleshchevo Lake National Park. *In: Ekologiya i ratsional'noe prirodopol'zovanie: materialy vserossiyskoy nauchno-prakticheskoy konferentsii* [Ecology and rational nature management: materials of the All-Russian scientific-practical conference (Yaroslavl, Russia, 12–16 September 2017)]. Yaroslavl – Periaslavl-Zaleskiy: 47–52 (in Russian).
- Waltl J. 1839. *Reise durch Tyrol, Oberitalien und Piemont nach dem südlichen Spanien, 2. Auflage.* Passau: Pustet'sche Buchhadlung. 247 + 120 p.
- Wanntorp H.-E. 2008. The identity of *Mantura obtusata* (Gyllenhal 1813) (Coleoptera, Chrysomelidae) – a Nordic-Central European mismatch. *Entomologisk Tidskrift*. 129(2): 99–101.
- Warchalowski A. 1998. Fauna Polski. T. 20. Cz. VI: Chrysomelidae. Stonkowate (Insecta: Coleoptera), Podrodzina Halticinae: rodzaje Hermaeophaga – Dibolia. Warszawa: Polskie towarzystwo entomologiczne. 360 p.
- Warchalowski A. 2010. The Palearctic Chrysomelidae. Identification keys. Vol. 1, 2. Warszawa: Natura Optima Dux Foundation. 1212 p.
- Weise J. 1886. Galerucinae. Lieferung 4. *In: Naturgeschichte der Insecten Deutschlands. Erste Abteilung Coleoptera. Sechster Band.* Berlin: Nicolaische Verlags-Buchhandlung: 569–768.
- Westwood J.O. 1840. An Introduction to the Modern Classification of Insects, Vol. 2. London: Longman, Orme, Brown, Green, and Longmans. 587 p.
- Zaitsev Yu.M., Medvedev L.N. 2009. Lichinki zhukov-listoedov Rossii [Larvae of leaf beetles of Russia]. Moscow: KMK Scientific Press Ltd. 246 p. (in Russian).
- Zhivotovskaya A.A. 1957. To the fauna of Halticinae (Chrysomelidae) of the Transcarpathian region. *Doklady i soobshcheniya Uzhgorodskogo gosudarstvennogo universiteta. Seriya biologicheskaya*. 1: 51–53 (in Russian).

Received / Поступила: 23.09.2020

Accepted / Принята: 12.11.2020

Published online / Опубликована онлайн: 22.12.2020